

## Lichens new for Slovakia collected in the National Park Muránska planina (W Carpathians)

ZDENĚK PALICE<sup>\*,#</sup>, ANNA GUTTOVÁ<sup>\*\*</sup> AND JOSEF P. HALDA<sup>\*\*\*</sup>

Palice Z., Guttová A. & Halda J. P. 2006. Lichens new for Slovakia collected in the National Park Muránska planina (W Carpathians). In: Lackovičová A., Guttová A., Lisická E. & Lizoň P. (eds.), Central European lichens – diversity and threat, p. 179 – 192. Mycotaxon Ltd., Ithaca.

**Abstract** – Seventeen new lichen taxa for Slovakia from the National Park Muránska planina (Central Slovakia, W Carpathians) are reported and shortly commented upon. They include both widely distributed, presumably undercollected, pioneer or ephemeral lichens (*Absconditella trivialis*, *Arthonia muscigena*, *Gyalidea diaphana*, *Micarea lithinella*, *Trapeliopsis glaucolepidea*, *Vezeada rheocarpa*) as well as lichenogeographically interesting records (*Buellia violaceofusca*, *Calicium montanum*, *C. pinastri*, *Caloplaca lucifuga*, *Chaenotheca sphaerocephala*, *Lecanora exspersa*, *Rinodina polysporoides*, *Verrucaria geophila*, *V. hegetschweileri*). *Protoparmelia hypotremella* and *Scoliosporum schadeanum* are probably partly undercollected species with a subatlantic bias and are apparently widespread in Slovakia, though not common.

**Key words** – lichenized fungi, diversity, threat, lichenogeography

### Introduction

The National Park Muránska planina (Central Slovakia, W Carpathians) is a sparsely inhabited and substantially forested area (81% of total area of the park, 65% of its protective zone; cf. Šmídt 2001). It shelters sensitive lichens of untouched natural habitats as well as of man-made/induced semi-natural habitats (road-ditches, alleys). The most spectacular natural ecosystems include exposed limestone/marble outcrops and cliffs forming ridges of the plateau (e.g. Poludnica) or topping the mountains Vrbiarka and Cigánka (hosting e.g. stenoendemic shrub *Daphne arbuscula* Čelak.). Important forest habitats cover old-growth deciduous forests – beech forests (e.g. Hrdzavá, Poludnica), oak forests on warmer slopes (Poludnica, Šiance and Cigánka), ravine forests with lime-trees, ashes and sycamores (Hrdzavá) and old-growth boreal forests on N slopes of Mt. Kľak. A raised peat-bog with *Pinus mugo* – a unique ecosystem in the National Park – occurs in the valley Hrdzavá dolina. The

\*Institute of Botany, Academy of Sciences of the Czech Republic, CZ-252 43 Průhonice, Czech Republic

\*\*Institute of Botany, Slovak Academy of Sciences, Dúbravská cesta 14, SK-845 23 Bratislava, Slovakia

\*\*\*Museum of Orlické hory, Jiráskova 2, CZ-516 01 Rychnov nad Kněžnou, Czech Republic

#Department of Botany, Faculty of Science, Charles University, Benátská 2, CZ-128 Prague 2, Czech Republic  
palice@ibot.cas.cz, anna.guttova@savba.sk, halda@jjh.cz

preservation of natural habitats is high, which is well documented by the relatively rich occurrence of the endangered epiphytic lichen *Lobaria pulmonaria* (Guttová & Palice 2001). The lichen flora of Muránska planina as a whole is exceptionally rich and unique vis-a-vis the above mentioned habitat richness and the fact that both montane and thermophilic species meet here on relatively small territory. The recent research focused on selected valuable small-scale areas in the SW part of the National Park and added 42 new species for Slovakia (Guttová & Palice 1999, 2002, 2004), i.e. species not included in the latest Checklist of Slovak lichens (Pišút et al. 1998, Bielczyk et al. 2004). Additional novelties originate either from other sites of the National Park or from already published inventories but were determined ex post after the results had been published.

## Materials and methods

Localities listed in the section Specimens examined are fully cited according to the labels. Altitude is given in meters above sea level. The specimens are deposited in PRA (followed by Z. Palice's collection number), SAV and private collections of J. P. Halda (hb. Halda) and Ch. Printzen (hb. Printzen). The acronyms used in the text follow Holmgren et al. (1990).

## Results

Taxa treated here represent both frequent and rare lichens. Some of the lichens are probably common, but so far overlooked in Slovakia and additional findings are expected. On the other hand there are also species of natural/relic habitats (i.e. lichens collected in oak forests, calcareous outcrops in open grasslands) with apparently restricted distribution in Slovakia. The species may be divided into six groups according to ecological/distributional characteristics:

- a) ephemeral and pioneer, widely distributed but undercollected species occurring in transient habitats (*Absconditella trivialis*, *Arthonia muscigena*, *Gyalidea diaphana*, *Micarea lithinella*, *Trapeliopsis glaucolepidea*, *Vezdaea rheocarpa*);
- b) boreal/montane species – occurring in peat-bogs and old-growth coniferous forest in the N part of Muránska planina (*Calicium pinastri*, *Chaenotheca sphaerocephala*, *Lecanora exspersa*);
- c) suboceanic-temperate old forest species (*Buellia violaceofusca*, *Caloplaca lucifuga*);
- d) southern/subatlantic taxa with submediterranean distribution not known from Northern Europe (*Calicium montanum*, *Rinodina polysporoides*, *Verrucaria geophila*);
- e) putatively suboceanic species (*Protoparmelia hypotremella*, *Scoliciosporum schadeanum*);

- f) poorly known taxa so far recorded only few times world-wide (*Verrucaria hegetschweileri*).

## The species

### *Absconditella trivialis* (Willey ex Tuck.) Vězda

Ephemeral lichen typically growing on sandy/loamy soil, e.g. at road-ditches. It is undercollected, however as it is an acidophilic species it is apparently missing in E part of the National Park which is a purely limestone area.

**SPECIMEN EXAMINED.** **Závadka nad Hronom.** the valley of Hronec, ca 300m W of Pätina game-keeper house (confluence with the Dudlavka), N 48°47'10-15", E 19°56', on sandy soil at road-side, alt. 810-820m, 5.V. 2001 Š. Bayerová, A. Guttová, J. Halda & Z. Palice (PRA 5309).

### *Arthonia muscigena* Th. Fr.

Small, apparently ephemeral lichen, which is widely overlooked. It occupies a wide range of substrates in moist, microclimatically suitable habitats. Recently Fryday (2004) forwarded nomenclature questions resulting from a thorough study of original material of *Lecidea lapidicola* Taylor, which is referable to *A. muscigena*.

**SPECIMENS EXAMINED.** **Zlatno.** The Havraník valley - a forest clearing, N 48°48'15-30", E 20°03'15-20", on decaying wood in the bedrock of the brook, *Micarea misella* associated, alt. 800-900m, 4.V. 2001 Š. Bayerová, J. Halda & Z. Palice (PRA 5471). **Muráň.** Javorníková dolina valley - the bottom (N 48°44', E 20°01'25"), on bark of *Fraxinus*, alt. 425-430m, 27.X. 2001 A. Guttová & Z. Palice (PRA 5564); *ibid.*, on bark of *Corylus*, alt. 460m, 8.V. 2001 A. Guttová, J. Halda & Z. Palice (PRA 5614). **Závadka nad Hronom.** The valley Za Nehovým, mouth of a side valley parallel with the valley Teplá dolina trending towards the cave Mochnatá (NNR Veľká Stožka), bark of *Acer pseudoplatanus* and decayed bryophytes growing on it, alt. ca 840m, 17.X. 2002 Š. Bayerová, A. Guttová & K. Kresáňová (SAV).

### *Buellia violaceofusca* Thor & Muhr

Described from S Sweden and only tentatively placed into the genus *Buellia* (Thor & Muhr 1991). Known only in sterile state having no secondary substances. Recognized by pale grey thallus and violet brown soralia. Characteristic for dry bark of old deciduous trees, especially *Quercus*. Reported also from Austria (Poelt 1994), Estonia (Thor & Nordin 1998), Belgium (Sérusiaux et al. 1999) and British Isles, where it is considered a Near Threatened species (Woods & Coppins 2003) as well as in Sweden (Gärdenfors 2005).

**SPECIMEN EXAMINED.** **Muráň.** Nature Reserve Poludnica, NNW-SSE oriented crest (N 48°45'40-45", E 20°02'03-05"), on bark of old *Quercus*, alt. 850m, 14.V. 2002 A. Guttová, J. Halda & Z. Palice (PRA 6264, cum *Caloplaca lucifuga*).

### *Calicium montanum* Tibell

The lichen superficially resembles a *Cyphelium* species, as the stalk is very short and hidden in the thick, well developed, glaucous thallus. It could be mistaken also for a short-stalked variety of *Calicium quercinum*, which however differs distinctly by its spiral ascospore ornamentation. In all cases it formed extensive patches on very hard oak wood. The species reaches northernmost European limit in the area, similarly like the epiphytes *Leptogium hildenbrandii* and *Physconia servitii* (Pišút 1990), saxicolous *Caloplaca adriatica* (Guttová & Palice 2004) or below treated *Verrucaria geophila*. *Calicium montanum* was recently described from mountains of southern Europe reaching northwards to Germany (Tibell 1999b) and Switzerland (Scheidtger et al. 2002). In Spain only four localities were recorded recently and it is regarded as an Endangered species there (Martínez et al. 2003). Known also from N Africa and Northern America (Kolb & Spribille 2001). Except *Quercus* it may also grow on hard wood of *Castanea* and *Pinus*.

**SPECIMENS EXAMINED.** Muráň. Mt. Šiance - S-SSE slope, light scree forest (N 48°46'10", E 20°04'30"), on hard wood of *Quercus*, alt. ca 860m, 15.V. 1998 Z. Palice, det. L. Tibell (PRA 327); *ibid.*, on hard wood of snag of *Quercus*, alt. 940-1000m, 7.V. 2001 Š. Bayerová, J. Halda & Z. Palice (PRA 5301); *ibid.*: a crest ENE of Šiance, N 48°46.45', E 20°05.62', well-lit deciduous forest (*Quercus*, *Tilia*, *Fraxinus*) on SSE facing slopes, on wood of branch of living oak, alt. 860m, 26.V. 2005 Z. Palice (PRA 9031); Nature Reserve Poludnica, NNW-SSE oriented crest (N 48°45'29", E 20°02'01"), on wood of *Quercus* snag, alt. 690-700m, 24.V. 2003 A. Guttová, J. Halda, Z. Palice, C. Printzen & I. Schmitt (hb. Halda 4814, 5670, PRA 6309, hb. Printzen).

### *Calicium pinastri* Tibell

This is a typical lichen of open boreal forest. It preferably grows on flaking bark of younger and middle-aged (faster growing) conifers, mostly pines, more rarely deciduous trees (*Betula*) or occasionally also on wood. Associated lichen taxa at the Slovak locality include other boreal-montane epiphytic/epixylic lichens, e.g. *Calicium glaucellum*, *C. trabinellum*, *Lecidea nylanderii* and *Pycnora sorophora*. *C. pinastri* was described from boreal forests in Finland, the Czech Republic and Germany (Tibell 1999) and later discovered also in Estonia (Jüriado et al. 2000), Sweden (Jonsson 2003) and Austria (Berger & Priemetzhofer 2005). One of the two Slovak specimens (hb. Palice 4044) was already published in a phylogenetic analysis on calicioid lichens based on ITS rDNA (Tibell 2003), where *Calicium pinastri* shows a close relationship to the superficially dissimilar *C. montanum*. No details are provided for the Slovak specimen and it was mistakenly indicated to originate from the Czech Republic.

**SPECIMENS EXAMINED.** Muráň. The Hrdzavá valley, peat-bog "V machoch", ca N 48°44'55"- 48°45', E 19°59'50", on bark of *Larix decidua*, alt. 650-700m, 18.X.1999 A. Guttová & Z. Palice (PRA 5071); *ibid.* 28.X. 2000 A. Guttová, J. Halda & Z. Palice, conf. L. Tibell (PRA 4044).

### *Caloplaca lucifuga* Thor

This sterile sorediate lichen is generally confined to dry fissures of old oaks both in wood- and parklands (Wirth 1995). It was rarely collected also on *Ulmus glabra* (Berger & Türk 1993), *Castanea* (Poelt 1994), *Aesculus hippocastanum* (Arup et al. 1997), *Tilia* (Türk & Wunder 1999), *Carpinus* (Fałtynowicz 2003) or *Acer pseudo-platanus* (Palice et al. 2003). Characterized by immersed thallus and well-delimited ochre concave soralia. It is often associated with calicioid lichens (Wirth 1995). There are scattered reports throughout Europe, most of them originate from southern Scandinavia – the northernmost distributional limit of *Quercus* – its main host tree. *Caloplaca lucifuga* is regarded an indicator of long forest continuity there (Arup et al. 1997). In Finland treated as a critically endangered species (Rassi et al. 2001). In Austria, Switzerland and Denmark it is ranked among endangered lichens (Türk & Hafellner 1999, Scheidegger et al. 2002, Søchting & Alstrup 2002). In Great Britain it is considered a Vulnerable (Woods & Coppins 2003) and a Near Threatened in Sweden (Gärdenfors 2005).

**SPECIMENS EXAMINED.** Muráň. Mt. Šiance - S-SW slope, light deciduous forest with *Quercus* predominating (N 48°46'00-05", E 20°04'20"), on dry bark of *Quercus*, alt. 940-1000m, 7.V. 2001 Š. Bayerová, J. Halda & Z. Palice (PRA 5305); Nature Reserve Poludnica, NNW-SSE oriented crest (N 48°45'40-45", E 20°02'03-05"), on bark of old *Quercus*, alt. 850m, 14.V. 2002 A. Guttová, J. Halda & Z. Palice (PRA 6264); *ibid.*, alt. 745m, 28.V. 2005 Z. Palice (PRA 8876).

### *Chaenotheca sphaerocephala* Nádv.

Described during second World War from Chile and known only from the type locality for a long time. Recently a number of collections from humid forests were attributed to this taxon, and it was subsequently reported from various sites of both hemispheres (Tibell 1998, 1999a, Palice 1999, Selva & Tibell 1999, Holien 2001, Titov 2001). In Europe, the species occurs mainly in natural old-growth boreal forests (Palice 1999, Tibell 1999a, Santesson et al. 2004). The Slovak specimen was collected in an old-growth humid stand on a steep slope. The ecology (bark of very shaded parts at very base of old spruce trees) fits well to previously reported records as its chemistry does. Finely granular thalli contain traces of obtusatic and barbatic acids, which corresponds with Tibell (1999a). These substances are produced also by *Ch. hygrophila* and *Ch. stemonea* or *Lepraria obtusatica* (Tibell 1999a, Tønsberg 1992), but the species feature different anatomical characters. In the field it could be mistaken for *Chaenotheca gracilenta* which may occur in similar habitats. These two species however are readily distinguishable by photobionts and ascospore size. According to Tibell (in litt.) the European specimens slightly differ from the type collection and both putative taxa are currently under taxonomic study. It is red-listed in Sweden as a Vulnerable species (Gärdenfors 2005).

**SPECIMEN EXAMINED.** Muráň. Mt. Klak [1409] - N facing steep slopes around Machnatá cave, ca 48°47' N, 19°58' E, at base of old *Picea*, alt. 1200-1300m, 5.V. 2000 A. Guttová, J. Halda, V. Orthová & Z. Palice (hb. Halda 4405, PRA 4669).

### *Gyalidea diaphana* (Nyl.) Vězda

This lichen is usually reported from natural sites on submerged stones in streams and humid places (e.g. Purvis et al. 1992, Santesson et al. 2004), however it also may occupy seminatural or anthropogenic habitats like e.g. mine quarries or road-ditches (Anonymus 2002), where it is growing on free-lying pebbles on the ground and accompanied e.g. by *Trapelia* sp. div. and *Micarea lithinella*. This is also the case of Slovak locality. Similar ecology was observed by the authors also in the Bohemian Forest and eastern Turkey.

**SPECIMEN EXAMINED.** Závadka nad Hronom. Fabova dolina valley, a managed spruce forest along forestry-trail traverse 0.75km NNW of the saddle “Burda”, 48°46'10" N, 19°54'10" E, on half-immersed siliceous pebbles at the trail-side, alt. 1100m, 5.V. 2001 Š. Bayerová, A. Guttová, J. Halda & Z. Palice (PRA 5470).

### *Lecanora exspersa* Nyl.

Rather poorly known epiphytic/epixylic sorediate species described from Retezat Mts. in Romania (Nylander 1875, Hazslinszky 1884; original material deposited in W); known from Montenegro (Vězda 2000), Switzerland (Clerc 2004), Austria (Wittmann & Türk 1989), Germany (Türk & Wunder 1991, Printzen et al. 2002) and Slovenia (Mrak et al. 2004). More localities are reported from Finland (cf. Santesson et al. 2004). Although red-listed, in the Alps it is not a rare lichen, ranked among species of Least concern in Switzerland (Scheidegger et al. 2002) and Vulnerable species in Austria (Türk & Hafellner 1999). In Muránska planina the extensive white-grey sorediate (not continuously) patches with dispersed, often pruinose apothecia covered almost whole the trunk of a young larch in raised-bog but not observed anywhere else around. Thallus reacts K+ yellow and contains atranorin and 1-2 unidentified fatty acids not belonging to roccellic acid (Tønssberg et al. 2001). Similar sorediate species occurring in Europe include *L. farinaria* Borrer and *L. mughosphagneti* Poelt & Vězda. Both species are usually sterile and their soralia often become soon confluent (Lumbsch et al. 1997, Tønssberg et al. 2001), and they contain atranorin like *L. exspersa*. In addition, *L. farinaria* contains roccellic acid (Tønssberg 1992, Tønssberg et al. 2001), while *L. mughosphagneti*, producing protocetraric acid, does not contain any fatty acid (Lumbsch et al. 1997). None of these species is known from Slovakia so far.

**SPECIMENS EXAMINED.** Muráň. The Hrdzavá valley - peat-bog “V machoch”, ca 48°44'55"- 48°45' N, 19°59'50" E, on bark of rather young *Larix decidua*, alt. 700-750m, 18.X.1999 A. Guttová & Z. Palice, conf. H. T. Lumbsch (PRA 2209, SAV); *ibid.*, 28.X. 2000 A. Guttová, J. Halda & Z. Palice (hb. Halda 4542, PRA 4051).

***Micarea lithinella* (Nyl.) Hedl.**

Widely distributed lichen, presumably nearly cosmopolitan. It grows as a pioneer on siliceous rocks, such as freshly exposed stones and small pebbles on the ground. In the area it occurs exclusively in W part of the national park which is siliceous. It is known also from NE Slovakia.

**SPECIMENS EXAMINED.** **Tisovec.** The Rimava valley, by the green-marked tourist footpath, ca 4,5km S of Mt. Fabova hoľa [1439], N 48°44', E 19°53'30", on silic. underhang together with *Micarea sylvicola*, alt. ca 750m, 13.V. 1998 Z. Palice (PRA 369); *ibid.*, N 48°44'00", E 19°53'50", on loose silic. at road-cutting, alt. ca. 600m, 13.V. 1998 Z. Palice (PRA 441); Mt. Fabova hoľa [1439] - SW slopes, pasture ca 0.5km ENE of the point 1092.4m (N 48°45'55", E 19°52'), on shaded side of fresh solitary silic. boulder, alt. ca 1100m, 14.V. 1998 Z. Palice (PRA 409). **Závodka nad Hronom.** Fabova dolina valley, a managed spruce forest along forestry-trail traverse 0,75km NNW of the saddle "Burda", N 48°46'10", E 19°54'10", on half-immersed siliceous pebbles at the trail-side, alt. 1100m, 5.V. 2001 Š. Bayerová, A. Guttová, J. Halda & Z. Palice (PRA 5307).

***Protoparmelia hypotremella* van Herk, Spier & V.Wirth**

The species may be overlooked for a stunted *Hypocenomyce caradocensis*, which is often found also in sterile state. In point of fact, *P. hypotremella* was not known fertile until report by Brodo & Aptroot (2005). The identity was usually revealed thanks to the parasymbiotic calicioid species *Sphinctrina anglica* Nyl. overgrowing its thallus. According to Aptroot et al. (1997) and Nordin & Hermansson (1999) this parasymbiont may grow on the thallus of both *P. hypotremella* and closely related *P. oleagina* (Harm.) Coppins. *P. hypotremella* differs from *P. oleagina* by its paler thallus (greyish to pale olivaceous to buff colour) consisting of dispersed isidia-like granules to squamules while *P. oleagina* has darker (brown to dark brown colour with greyish to olivaceous hue), more continuous thallus and true isidia. Both species contain lobaric acid, however only *P. hypotremella* is UV+ (Aptroot et al. 1997, Nordin & Hermansson 1999, Brodo & Aptroot 2005). It could be confused also with *P. ochrocoeca*, which unlike *P. hypotremella* produces round to hemispherical, pale to dark reddish brown squamules, lobaric acid is absent. It is not parasitized by *Sphinctrina*. Slovak material shows a distinct UV+ positive reaction and matches well the descriptions of *P. hypotremella* by the authors cited above.

**SPECIMENS EXAMINED.** **Muráň.** Nature Reserve Poludnica, NNW-SSE oriented crest (N 48°45'36", E 20°02'01"), on wood of bole of *Quercus* snag, alt. 790m, 24.V. 2003 A. Guttová, J. Halda, Z. Palice, C. Printzen & I. Schmitt (hb. Printzen, PRA 6298, SAV – cum *Sphinctrina anglica*).

***Rinodina polysporoides* Giralt & H.Mayrhofer**

This is a species distributed in mountains of Mediterranean reaching Central Europe (Giralt & Mayrhofer 1994), where it is known so far from Slovenia, Switzerland and Austria. Both in Austria and Switzerland it is treated as a Vulnerable

species (Türk & Hafellner 1999, Scheidegger et al. 2002). The presented collections represent the northernmost known localities of the taxon. Another species of *Rinodina* with polysporic asci occurring in Central Europe – *R. polyspora* – is a boreal-montane lichen with *Physcia*-type ascospores while *R. polysporoides* has the *Dirinaria*-type of ascospores.

**SPECIMENS EXAMINED.** **Muráň.** W-slope of Žabica hill (1km SW of Muráň), a row of middle-aged *Juglans*-trees on a pasture, on bark of solitary *Juglans*, alt. 470m, 12.V. 1999 A. Guttová, J. Halda & Z. Palice, conf. H. Mayrhofer (PRA 2076); *ibid.* 29.X.2000 A. Guttová & Z. Palice (PRA 4023); “Na Kopanke” - a walnut orchard near forest margin (Poludnica reserve) ca 2km NW of Muráň village, SE foot of the point Vyšný Kostolec [1144,3], N 48°45.02', E 20°01.73', on twig of *Juglans*, alt. 440-450m, 28.V.2005 Z. Palice (PRA 8934).

### *Scoliciosporum schadeanum* (Erichs.) Vězda

The species is doubtfully distinct from the later described but much frequently recorded *S. pruinosum* (P. James) Vězda, which allegedly differs in (richer) development of epipsamma in hymenium/exciple and by more richly branched hamathecium. The species has subatlantic distribution in Europe. Although it is easily overlooked the species may well be spreading during last decades as indicated by more recent finds in different parts of the Czech Republic (Palice et al. 2003).

**SPECIMENS EXAMINED.** **Zlatno.** 2km S of Zlatno, deciduous forest on bank of Havraník brook, on bark of *Acer pseudoplatanus*, alt 900-950m, 10.V. 1999 J. Halda & Z. Palice (hb. Halda 3221). **Muráň.** At forest-roadside on tourist red-marked footpath, ca 4km N of Muráň village (N 48°46'30", E 20°03'20"), on bark of *Fagus*, alt. 950m, 11.V. 1998 Z. Palice (PRA 299). **Závadka nad Hronom.** The valley of Hronec (Fabova dolina valley) - the upper part, N 48°46'40", E 19°54'30", on bark of *Acer pseudoplatanus*, alt. 1000m, 5.V. 2001 Š. Bayerová, A. Guttová, J. Halda & Z. Palice (PRA 5462).

### *Trapeliopsis glaucolepidea* (Nyl.) G.Schneider

The collected specimen represents a diminished lignicolous form sometimes called *Trapeliopsis percrenata*. It is apparently a widespread lichen in humid regions of Slovakia. In Central Europe it favours peaty soils, humus and decaying wood in humid forests and may be overlooked for *Cladonia* squamules as the apothecia are only rarely developed. The species contains an UV+ blue-white unknown substance. It usually forms fine pale-creamy, on exposed sites bluish grey orbicular soralia which later merge to form marginal soralia. It may readily be distinguished from *Cladonia* squamules by a transient KC+ red spot reaction.

**SPECIMEN EXAMINED.** **Muráň.** Nature Reserve Veľká Stožka, Mt. Kľak [1409] - N facing slopes above Machnatá cave, N 48°46.92', E 19°57.69', on decaying wood (vertical part) of (?) *Picea* stump, alt. 1305-1310m, 27.VIII.2005 Z. Palice (PRA 8910).



***Verrucaria geophila* Zahlbr.**

A terricolous species with most localities in the Mediterranean and Balkan. So far the northernmost record was from Austria (Wittmann & Türk 1994). It is characterized by relatively large perithecia (ca 0.35–0.5 mm) and ascospores (24–33 × 9.5–13 µm), and a well developed areolate/subsquamulose glossy olivaceous grey/green thallus.

**SPECIMENS EXAMINED.** Muránska Huta. Valley of Trsteník, by road-side, over mosses on ground, alt. 900m, 23.IX.1995 Z. Palice, det. O. Breuss (PRA s.n.). Muráň. Mt. Vrbiarka - SE exposed rocks above Javorníková valley (ca. N 48°44'30", E 20°00'), on humus and plant debris at rather steep calcareous rock, alt. 1000-1050m, 27.X.2000 A. Guttová, J. Halda & Z. Palice (PRA 4704).

***Verrucaria hegetschweileri* Körber**

A poorly known corticolous species of *Verrucaria* described from Switzerland, and afterwards recorded only from the Czech Republic (Breuss 1998) and Belgium (Sérusiaux et al. 2003).

**SPECIMEN EXAMINED.** Zlatno. Valley of Sviniarka brook, nature reserve "Zlatnica", N 48°49'26", E 20°05'41", on weathered bark of old *Fagus*, alt. 795m, 25.X.2001 Z. Palice, det. O. Breuss (PRA 5749).

***Veizdaea rheocarpa* Poelt & Döbbeler**

In the area of Muránska planina 4 species of the genus *Veizdaea* were collected so far (*V. aestivalis*, *V. retigera*, *V. rheocarpa*, *V. stipitata*). All the species are minute, easily overlooked ephemeral lichens sharing a similar ecology. *V. rheocarpa* is reported here as new to Slovakia. It is distinguished from related species by microscopic morphological characters of the thallus and apothecia (see Purvis et al. 1992).

**SPECIMENS EXAMINED.** Muráň. Mt. Veľký Cigán - SE slope, at yellow tourist footpath, ca 0.4-0.5km E of the point 1218m, N 48°46'55", E 20°01'40", on bare sandy soil among roots of an eradicated tree, alt. 1120-1130m, 11.V.1998 V. Orthová & Z. Palice (PRA 417); By red tourist footpath ca 0,5km W of abandoned gamekeeper-house "Maretkino", ca N 48°46', E 20°01', on detritus at forest-roadside, alt. 980m, 12.V.1997 A. Guttová & Z. Palice (PRA 1753).

## Acknowledgements

It is our great pleasure to dedicate this paper to Ivan Pišút who contributed so much to the understanding of lichens in Slovakia, Mediterranean and Asia and continuously encourages younger lichenologists. Our dear colleagues Štěpánka Slavíková-Bayerová (Průhonice), Katarína Kresáňová (Bratislava), Viera Slezáková (Bratislava), Christian Printzen (Frankfurt am Main) and Imke Schmitt (Chicago) attended and helped us to collect the material in the field. We are grateful to Štěpánka Slavíková-Bayerová also for TLC of *Ch. sphaerocephala*. Othmar Breuss

(Wien) kindly determined and verified most of *Verrucaria* specimens. The identity of *Lecanora exspersa* and *Rinodina polysporoides* was confirmed by Thorsten Lumbsch (Chicago) and Helmut Mayrhofer (Graz), respectively. Leif Tibell (Uppsala) kindly revised several *Calicium* specimens and reviewed the manuscript. Drahoš Blanár (Revúca) helped us many times in various unexpected ways during the field research. ZP acknowledges continued support by Grant Agency of Academy of Sciences of the Czech Republic (projects AV0Z60050516) and by the Ministry of Education, Youth and Sports, Czech Republic, project no. 21620828. AG is obliged to the support by Science and Technology Assistance Agency under the contract No. APVT-51-005102.

### Literature cited

#### ANONYMUS

2002 Fundmeldungen. Akt. Lichenol. Mitt. NF 9: 28-29.

#### APTROOT A., DIEDERICH P., VAN HERK C. M., SPIER L. & WIRTH V.

1997 *Protoparmelia hypotremella*, a new sterile corticolous species from Europe, and its lichenicolous fungi. Lichenologist 29: 415-424.

#### ARUP U., EKMAN S., KÄRNEFELT I. & MATTSSON J-E. (EDS.)

1997 Skyddsvärda lavar i sydvästra Sverige. SBF-förlaget, Lund.

#### BERGER F. & PRIEMETZHOFFER F.

2005 Neue und bemerkenswerte Funde von Flechten aus Oberösterreich Österreich. Beitr. Naturk. Oberösterreichs 14: 3-18.

#### BERGER F. & TÜRK R.

1993 Neue und seltene Flechten und lichenicole Pilze aus Oberösterreich. Österreich. Linzer Biol. Beitr. 25: 167-204.

#### BIELCZYK U., LACKOVIČOVÁ A., FARKAS E., LÓKÖS L., LIŠKA J., BREUSS O. & KONDRATYUK S. YA.

2004 Checklist of lichens of the Western Carpathians. Biodiversity of the Carpathians 1: 5-181. W. Szafer Inst. Bot., Pol. Acad. Sci., Kraków.

#### BREUSS O.

1998 Drei neue holz- und borkenbewohnende *Verrucaria*-Arten mit einem Schlüssel der bisher bekannten Taxa. Linzer Biol. Beitr. 30: 831-836.

#### BRODO I. M. & APTROOT A.

(2005) Corticolous species of *Protoparmelia* (lichenized Ascomycotina) in North America. Can. J. Bot. 83: 1075-1081.

#### CLERC P.

2004 Les champignons lichénisés de Suisse. Catalogue bibliographique complété par des données sur la distribution et l'écologie des espèces. Cryptog. Helv. 19: 1-320.

#### DIEDERICH P.

2003 New species and new records of American lichenicolous fungi. Herzogia 16: 41-90.

- FAŁTYNOWICZ W.  
2003 The lichens, lichenicolous and allied fungi of Poland - an annotated checklist. W. Szafer Inst. Bot., Pol. Acad. Sci., Kraków.
- GÄRDENFORS U. (ED.)  
2005 Rödlistade arter i Sverige 2005. ArtDatabanken, SLU, Uppsala.
- GIRALT M. & MAYRHOFER H.  
1994 Four corticolous species of the genus *Rinodina* (lichenized Ascomycetes, *Physciaceae*) with polyspored asci. *Herzogia* 10: 29-37.
- GUTTOVÁ A. & PALICE Z.  
1999 Lišajníky Národného parku Muránska planina I - Hrdzavá dolina. In: UHRIN M. (ed.), Výskum a ochrana prírody Muránskej planiny 2. pp. 35-47. MŽP SR Bratislava a Správa NP Muránska planina, Revúca.  
2001 Výskyt jamkatca pľúcneho (*Lobaria pulmonaria*) v NP Muránska planina (Stredné Slovensko). *Bull. Slov. Bot. Spoločn.*, Bratislava, 23: 39-43.  
2002 Lišajníky Národného parku Muránska planina II - Javorníková dolina. Výskum a ochrana prírody Muránskej planiny 3: 53-68.  
2004 Lišajníky Národného parku Muránska planina III - Cigánka. *Reussia* 1, Suppl. 1/2004: 5-40.
- FRYDAY A.  
2004 A New Species of *Fuscopannaria* with a Green Photobiont, and Other Taxonomic Innovations and New Records of Lichenized-Fungi from Alaska. *Bryologist* 107: 173-179.
- HAZSLINSZKY F.  
1884 A Magyar birodalom zuzmó-flórája. Budapest.
- HOLIEN H.  
2001 Additions to the Norwegian flora of lichens and lichenicolous fungi II - with some further distributional notes on Norwegian Caliciales. *Graphis Scripta* 12: 51-58.
- HOLMGREN P. K., HOLMGREN N. H. & BARNET L.  
1990 Index herbariorum. Part I: The herbaria of the world. 86th ed. Bronx, New York Botanical Garden.
- JONSSON F.  
2003 *Calicium pinastri* new to Sweden. *Graphis Scripta* 14: 5-6.
- JÜRIADO I., LÖHMUS P. & SAAG L.  
2000 Supplement to the second checklist of lichenized, lichenicolous and allied fungi of Estonia. *Folia Cryptog. Estonica* 37: 21-26.
- KOLB A. & SPRIBILLE T.  
2001 *Calicium corymellum* (Ach.) Ach. in the United States, and *Calicium montanum* Tibell new for North America. *Evansia* 18: 90-92.
- LUMBSCH H. T., PLÜMPER M., GUDERLEY R. & FEIGE G. B.  
1997 The corticolous species of *Lecanora* sensu stricto with pruinose apothecial discs. *Symb. Bot. Ups.* 82/1: 131-162.
- MARTÍNEZ I., ARAGÓN G., SARRIÓN F. J., ESCUDERO A., BURGAS A. R. & COPPINS B. J.  
2003 Threatened lichens in central Spain (saxicolous species excluded). *Cryptog. Mycol.* 24: 73-97.

- MRÁK T., MAYRHOFER H. & BATIČ F.  
 2004 Contributions to the lichen flora of Slovenia XI. Lichens from the vicinity of Lake Bohinj (Julian Alps). *Herzogia* 17: 107-127.
- NORDIN A. & HERMANSSON J.  
 1999 Floristic news from Sweden, Norway and Finland. *Graphis Scripta* 10: 13-20.
- NYLANDER W.  
 1875 Addenda nova ad Lichenographiam europaeam. *Continuatio tertia et vicesima* (23). *Flora* 58: 440-448.
- PALICE Z.  
 1999 New and noteworthy records of lichens in the Czech Republic. *Preslia* 71: 289-336.
- PALICE Z. ET AL.  
 2003 Lišejníky zaznamenané během 9. jarního setkání bryologicko-lichenologické sekce v Hajnici u Mirochova (CHKO Třeboňsko, 11.-14.4.2002). *Bryonora* 32: 7-17.
- PIŠŮT I.  
 1990 Nachträge zur Kenntnis der Flechten der Slowakei 12. *Zborn. Slov. Nár. Múz., Přír. Vedy, Bratislava*, 36: 9-13.
- PIŠŮT I., GUTTOVÁ A., LACKOVIČOVÁ A. & LISICKÁ E.  
 1998 Lichenizované huby (lišajníky). In: MARHOLD K. & HINDÁK F. (eds.), *Zoznam nižších a vyšších rastlín Slovenska*. p. 229-295. Veda, Bratislava.
- POELT J.  
 1994 Bemerkenswerte Flechten aus Österreich, insbesondere der Steiermark. *Mitt. Naturwiss. Ver. Steiermark* 124: 91-111.
- PRINTZEN C., HALDA J., PALICE Z. & TØNSBERG T.  
 2002 New and interesting lichen records from old-growth forest stands in the German National Park Bayerischer Wald. *Nova Hedwigia* 74: 25-49.
- PURVIS O. W., COPPINS B. J., HAWKSWORTH D. L., JAMES P. W. & MOORE D. M. (EDS.)  
 1992 The lichen flora of Great Britain and Ireland. Natural History Museum, London.
- RASSI P., ALANEN A., KANERVA T. & MANNERKOSKI I. (EDS.)  
 2001 The 2000 Red List of Finnish species. Helsinki, Ministry of the Environment & Finnish Environment Institute. [English Summary]
- SANTESSON R., MOBERG R., NORDIN A., TØNSBERG T. & VITIKAINEN O.  
 2004 Lichen-forming and lichenicolous fungi of Fennoscandia. Museum of Evolution, Uppsala.
- SCHEIDEGGER C. ET AL.  
 2002 Rote Liste der gefährdeten Arten der Schweiz: baum- und erdbewohnende Flechten. BUWAL-Reihe Vollzug Umwelt, Bern, Birmensdorf & Genève.
- SELVA S. B. & TIBELL L.  
 1999 Lichenized and non-lichenized calicioid fungi from North America. *Bryologist* 102: 377-397.
- SÉRUSIAUX E., DIEDERICH P., BRAND A. M. & VAN DEN BOOM P. P. G.  
 1999 New or interesting lichens and lichenicolous fungi from Belgium and Luxembourg. VIII. *Lejeunia* 162: 1-95.

- SÉRUSIAUX E., DIEDERICH P., ERTZ D. & VAN DEN BOOM P. P. G.  
2003 New or interesting lichens and lichenicolous fungi from Belgium, Luxembourg and northern France. IX. *Lejeunia* 173: 1-48.
- SØCHTING U. & ALSTRUP V.  
2002 Danish Lichen Checklist. Botanical Institute, University of Copenhagen, Copenhagen.
- ŠMÍD J.  
2001 Lesné hospodárstvo. In: UHRIN M. (ed.), Ročenka Správy NP Muránska planina. p. 57-58. Revúca.
- THOR G. & MUHR L-E.  
1991 *Buellia violaceofusca*, a new lichen from Sweden. *Lichenologist* 23: 11-13.
- THOR G. & NORDIN A.  
1998 16 lichens new to Estonia. *Folia Crypt. Estonica* 32: 123-125.
- TIBELL L.  
1998 Crustose mazaediate lichens and the *Mycocaliciaceae* in temperate South America. *Bibl. Lichenol.* 71: 1-107.  
1999a Calicioid lichens and fungi. In: AHTI T. et al (eds.), *Nordic Lichen Flora*. Volume 1. pp. 20-93. Nordic Lichen Society, Uddevalla.  
1999b Two new species of *Calicium* from Europe. *Mycotaxon* 70: 431-443.  
2003 *Tholurna dissimilis* and generic delimitations in *Caliciaceae* inferred from nuclear ITS and LSU rDNA phylogenies (*Lecanorales*, lichenized ascomycetes). *Mycol. Res.* 107: 1403-1418.
- TITOV A.  
2001 Further notes on calicioid lichens and fungi from the Gongga Mountains (Sichuan, China). *Lichenologist* 33: 303-314.
- TØNSBERG T.  
1992 The sorediate and isidiate, corticolous, crustose lichens in Norway. *Sommerfeltia* 14: 1-331.
- TØNSBERG T., TÜRK R. & HOFMANN P.  
2001 Notes on the lichen flora of Tyrol (Austria). *Nova Hedwigia* 72: 487-497.
- TÜRK R. & HAFELLNER J.  
1999 Rote Liste gefährdeter Flechten (Lichenes) Österreichs. 2. Fassung. In: NIKLFELD H. et al. (eds.), Rote Listen gefährdeter Pflanzen Österreichs. 2. Auflage. Grüne Reihe des Bundesministeriums für Umwelt, Jugend und Familie, Band 10. p. 187-228. Austria Medien Service, Graz.
- TÜRK R. & WUNDER H.  
1991 Die Kartierung der epiphytischen und epixylen Flechten im Nationalpark Berchtesgaden und dessen Vorfeld. *Schriftenreihe Bayer. Landesamt für Umweltschutz* 102: 79-91.  
1999 Die Flechten des Nationalparks Berchtesgaden und angrenzender Gebiete. Berchtesgaden, Nationalpark Berchtesgaden, Forschungsbericht 42, Nationalparkverwaltung Berchtesgaden.
- VÉZDA A.  
2000 Lichenes Rariores Exsiccati. Fasciculus 45 (numeris 441-450). Brno.

WIRTH V.

1995. Die Flechten Baden-Württembergs I, II. Eugen Ulmer Verlag, Stuttgart.

WITTMANN H. & TÜRK R.

1989 Zur Kenntnis der Flechten und flechtenbewohnenden Pilze von Oberösterreich und Salzburg II. *Herzogia* 8: 187-205.

1994 Flechten und Flechtenparasiten der Ostalpen II. *Ber. Bayer. Bot. Ges.* 64: 189-204.

WOODS R. G. & COPPINS B. J.

2003 A Conservation Evaluation of British Lichens. British Lichen Society, London.