
MYCOTAXON

ISSN (print) 0093-4666 (online) 2154-8889 Mycotaxon, Ltd. ©2018

January–March 2018—Volume 133, pp. 55–61

<https://doi.org/10.5248/133.55>

Four new records of *Leptogium* from China

YU XIA, LI-LI ZHAO, QING-FENG WU, HUA-JIE LIU*

College of Life Sciences, Hebei University, Baoding, 071002, China

* CORRESPONDENCE TO: liuhuajie@foxmail.com

ABSTRACT—Four *Leptogium* species are reported as new to China: *L. askotense*, *L. coralloideum*, *L. milligranum*, and *L. subtile*. Descriptions, illustrations, and comments are presented for the four species.

KEY WORDS—*Ascomycota*, *Peltigerales*, *Collembataceae*, lichen, taxonomy

Introduction

Forty-two species of the lichen genus *Leptogium* (Ach.) Gray have been reported from China (Wei 1991; Cao & al. 2012; Liu & al. 2010, 2012, 2013, 2014, 2015; Liu & Guan 2012; Wang & al. 2010; Xi & Liu 2014). Here we report *Leptogium askotense*, *L. coralloideum*, *L. milligranum*, and *L. subtile* as new to China. Although the genus was recently re-circumscribed (Otálora & al. 2014, Jaklitsch & al. 2016), we followed the traditional genus concept (Lumbsch & Huhndorf 2007, Kirk & al. 2008) in the present study.

Materials & methods

Morphological and anatomical studies were conducted following Liu & al. (2015). Specimens are deposited in the Herbarium Mycologicum Academiae Sinicae–Lichenes, Beijing, China (HMAS-L) and the Herbarium, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming, China (KUN).

Taxonomy

Leptogium askotense D.D. Awasthi, Norweg. J. Bot. 24: 63 (1977). PL. 1A

THALLUS foliose, orbicular or irregular in outline, 4–12 cm in diam.; UPPER SURFACE gray to brown when dry, glabrous, heavily wrinkled, dull or slightly

shiny near margin; LOBES round, 3–10 mm wide, horizontal, entire; ISIDIA and LOBULES absent; LOWER SURFACE paler, densely hairy; HAIRS white, 4–7 μm diam., >100 μm long, composed of a row of cylindrical cells. APOTHECIA abundant, laminal to sometimes submarginal, stalked; DISC red-brown, 2–4 mm in diam., smooth, dull, plane, concave to convex; STALK commonly tubular, not swollen, often with wrinkles and white tomenta; THALLINE MARGIN distinct, concolorous with the thallus, often with tomenta and wrinkles similar to stalk.

THALLUS homoiomorous, 100–250 μm thick; CORTEX on both sides consisting of a single layer of subglobose to irregular, thin-walled cells, 5–10 μm thick; PHOTOBIONT *Nostoc* in chains, 4–7 μm in diam., embedded in a gelatinous layer.

APOTHECIA zeorine, 400–500 μm thick; THALLINE EXCIPLE 100–120 μm thick laterally, with a cortex of a single layer of thin-walled cells; PROPER EXCIPLE euparaplectenchymatous, 50–80 μm thick, consisting of 5–8 layers of subglobose to irregular thin-walled cells; EPIHYMENIUM yellowish, 10–20 μm thick; HYMENIUM 120–150 μm thick; SUBHYMENIUM 40–70 μm thick; ASCI clavate, 8-spored; ASCOSPORES monostichous to distichous, ellipsoid, hyaline, muriform, 5–6-septate transversely, 1–2-septate longitudinally, (15)25–40 \times 10–15 μm , obtuse or acute at both ends.

Corticolous.

SELECTED SPECIMENS EXAMINED: CHINA. SICHUAN PROVINCE, Mt. Gonggashan, 29°37'N 101°53'E, alt. 2100 m, on bark, 23/VI/1982, X.Y. Wang & al. 8829 (HMAS-L 037125). YUNNAN PROVINCE, Lijiang County, Mt. Yulongshan, 27°10'N 100°18'E, alt. 3500 m, on bark, 25/I/2004, L.S. Wang & J. Wang 04-23160 (KUN). XIZANG PROVINCE, Zayü County, Mt. Songtaxueshan, 27°43'N 98°39'E, alt. 3400 m, on bark, 26/VI/1982, J.J. Su 2010 (HMAS-L 031687).

REMARKS: *Leptogium askotense* is characterized by the foliose thallus, the wrinkled upper surface, the tomentose and wrinkled thalline margin and stalks, and the euparaplectenchymatous proper exciple. It is reported from Himalayas of India (Awasthi & Akhtar 1977). Our specimens are morphologically similar to those reported from India and extend the species' range to SW China.

Leptogium trichophoroides P.M. Jørg. & A.K. Wallace resembles *L. askotense* in the tomentose, wrinkled thalline margin and the tomentose apothecial stalks, but differs by the darker upper surface and smaller spores (20–25 \times 10–12 μm , Cao & al. 2012; 19–22 \times 9–10 μm , Jørgensen 1997). *Leptogium delavayi* Hue is similar to *L. askotense* but has glabrous, swollen stalks and a glabrous thalline margin. *Leptogium arisanense* Asahina also resembles *L. askotense* but has sessile apothecia (Asahina 1936, Awasthi & Akhtar 1977, Wang & al. 2010).

Leptogium coralloideum (Meyen & Flot.) Vain.,
Ann. Acad. Sci. Fenn., Ser. A, 6(7): 110 (1915).

PL. 1B,C

THALLUS foliose, irregular in outline, 6 cm in diam.; UPPER SURFACE grayish blue when dry, glabrous, often striate and ridged, dull, isidiate; LOBES round, horizontal near center, somewhat erect near margin, <5 mm wide, the margins entire to somewhat lacerate; ISIDIA laminal, mostly on ridges, concolorous with the thallus or darker, not wrinkled, not apically collapsed, granular when young, cylindrical and coralloid when mature; LOWER SURFACE paler, irregularly striate, with scattered tufts of white hairs.

THALLUS homoiomerous, 100–200 µm thick; CORTEX on both sides consisting of a single layer of irregular, thin-walled cells of 5–7 µm in diam.; PHOTOBIONT *Nostoc* cells in chains, embedded in a gelatinous layer.

APOTHECIA not seen.

On mosses.

SPECIMEN EXAMINED: CHINA. HUNAN PROVINCE, Sangzhi County, Mt. Tianpingshan, 29°44'N 110°00'E, on moss, 19/VIII/1997, J.B. Chen & D.P. Wang 9606 (HMAS-L 031566).

REMARKS: *Leptogium coralloideum* is characterized by the coralloid isidia distributed mainly along ridges on upper surface, and the lower surface with striates and few hairs. It is widely distributed worldwide, especially common in tropical and temperate regions (Aragon & al. 2005, Jørgensen & Nash 2004, Jørgensen 1994). The single specimen from China is morphologically similar to those reported outside the country.

Five coralloid–isidiate species—*Leptogium burnetiae* C.W. Dodge, *L. cyanescens* (Rabenh.) Körb., *L. laceroides* B. de Lesd., *L. lichenoides* (L.) Zahlbr. [≡ *Scytinium lichenoides* (L.) Otálora & al.], and *L. pseudopapillosum* P.M. Jørg.—have been reported in China (Jatta 1902, Wei 1991, Jørgensen 1997, Aptroot & al. 2002, Wang & al. 2010). Both *L. burnetiae* and *L. pseudopapillosum* differ from *L. coralloideum* by the dense, >100 µm long hairs covering the lower surface; *L. lichenoides* differs by its semi-erect to erect lobes and marginal isidia; *L. cyanescens* differs by its thinner lobes and upper surface without ridges; and *L. laceroides* differs by its smooth upper surface and dense hairs on lower surface.

Leptogium milligranum Sierk, Bryologist 67: 290 (1964).

PL. 1D

THALLUS foliose, irregular in outline, 4 cm in diam.; UPPER SURFACE gray-green when dry, glabrous, longitudinally wrinkled, isidiate; LOBES irregular or somewhat indistinct, 2–8 mm wide, horizontal, often anastomosing, partly fenestrate, margin incised, occasionally with granular isidia; ISIDIA

laminal, numerous, granular, simple, sometimes also cylindrical and slightly branched, glabrous, without wrinkles; LOWER SURFACE paler, with scattered tufts of white hairs.

THALLUS homoiomerous, 180–280 µm thick; CORTEX on both sides consisting of a single layer of subglobose, thin-walled cells of 4–7 µm in diam.; PHOTOBIONT *Nostoc* cells in chains, embedded in a gelatinous layer.

APOTHECIA not seen.

Corticolous.

SPECIMEN EXAMINED: CHINA. YUNNAN PROVINCE, Yimen County, 24°40'N 102°06'E, alt. 1980 m, on bark, 24/IX/2012, L.S. Wang & al. 12-35195 (KUN).


REMARKS: *Leptogium milligranum* is characterized by the green thallus, numerous granular isidia and wrinkles on upper surface, the indistinct, anastomosing lobes, and the sparsely rhizinate lower surface. Four granular–isidiate species—*L. asiaticum* P.M. Jørg., *L. papillosum* (B. de Lesd.) C.W. Dodge, *L. saturninum* (Dicks.) Nyl., and *L. weii* H.J. Liu & S. Guan—have been reported in China (Jatta 1902, Zahlbruckner 1930, Wang & al. 2010, Liu & Guan 2012), but they have distinct, round, non-anastomosing lobes.

Leptogium milligranum has been recorded from North America, Central America, Galápagos Islands, and India (Sierk 1964, Awasthi & Akhtar 1979, Jørgensen & Nash 2004, Bungartz 2008). Our specimen falls within the range of morphological variation of those reported from outside China.

Leptogium subtile (Schrad.) Torss., Enum. Lich. Byssac. Scandin.: 54 (1843). PL. 1E,F

THALLUS minutely foliose, partly forming a pulvinate cushion, irregularly in outline, <4 cm in diam.; UPPER SURFACE bluish when dry, glabrous, smooth, dull; LOBES dense, membranaceous, deeply dissected, 1–2 mm wide near center, <0.5 mm wide near margin, free to suberect at margin, entire; ISIDIA and LOBULES absent; LOWER SURFACE concolorous with upper surface, without wrinkles, with scattered tufts of white hairs. APOTHECIA scattered, laminal, sessile; disc red brown, 0.2–0.4 mm in diam., dull; THALLINE MARGIN persistent, concolorous with thallus or paler, glabrous, entire, without wrinkles and isidia.

PLATE 1. *Leptogium askotense* (KUN Wang 04-23160). A. Thallus with apothecia. *Leptogium coralloideum* (HMAS-L 031566). B. Thallus; C. Upper surface, showing isidia on ridges. *Leptogium milligranum* (KUN Wang 12-35195). D. Thallus, showing dense, granular isidia. *Leptogium subtile* (KUN Wang 05-24245). E. Thallus; F. Lobes and apothecia. Scale bars: A = 20 mm; B, D, E = 10 mm; C = 5 mm; F = 2 mm.



THALLUS homoiomerous, 40–50 µm thick; CORTEX on both sides consisting of a single layer of subglobose to slightly irregular, thin-walled cells of 3–6 µm in diam.; PHOTOBIONT *Nostoc* in chains. APOTHECIA lecanorine; EPITHECIUM yellowish to sometimes brown, about 10 µm thick; HYMENIUM 100–150 µm thick; SUBHYMENIUM 20–30 µm thick; PROPER EXCIPLE not seen; ASCI clavate, 8-spored; ASCOSPORES colourless, ellipsoid, obtuse at both ends, muriform, 4–6-septate transversely, 1–2-septate longitudinally, 20–30 × 10–12 µm.

On moss-covered barks.

SPECIMEN EXAMINED: CHINA. YUNNAN PROVINCE, Yuanyang County, Huangcaoling Town, 22°57'N 102°37'E, alt. 1200 m, on bark, 16/1/2005, L.S. Wang 05-24245 (KUN).

REMARKS: *Leptogium subtile* is easily distinguished from other species in China by its minutely foliose and thin thallus, dissected, narrow and membranaceous lobes, smooth upper surface, and small, sessile apothecia. It was previously known from North America, Europe, and Asia (Jørgensen 1994, Kashiwadani & al. 2002, Jørgensen & Nash 2004, Aragón & al. 2005). It is rare in China with only one collection, which falls in the range of morphological variation of those from outside China.

Acknowledgements

This study was supported by the National Natural Science Foundation of China (31093440), Natural Science Foundation of Hebei Province (C2014201032), the Key Bioengineering Discipline of Hebei Province (1050-5030023), and the Key Laboratory of Microbial Diversity Research and Application of Hebei Province. The authors are grateful to Prof. Li-Song Wang (Kunming Institute of Botany, CAS), Prof. Jiang-Chun Wei and Ms. Hong Deng (Institute of Microbiology, CAS) for sending the specimens on loan, and Dr. Xin-Yu Wang (Kunming Institute of Botany, CAS) and Dr. Ze-Feng Jia (College of Life Sciences, Liaocheng University, China) for reading and improving the manuscript as presubmission reviewers.

Literature cited

- Aptroot A, Sparrius LB, Lai MJ. 2002. New Taiwan macrolichens. *Mycotaxon* 84: 281–292.
- Aragón G, Otálora MAG, Martínez I. 2005. New data on the genus *Leptogium* (lichenized *Ascomycetes*) in the Iberian Peninsula. *Nova Hedwigia* 80(1–2): 199–226. <https://doi.org/10.1127/0029-5035/2005/0080-0199>
- Asahina Y. 1936. Zwei neue arten von *Leptogium* aus Formosa. *Journal of Japanese Botany* 12(4): 250–255.
- Awasthi DD, Akhtar P. 1977. The genus *Leptogium* (sect. *Mallotium*) in India. *Norwegian Journal of Botany* 24(2): 59–71.
- Awasthi DD, Akhtar P. 1979. The lichen genus *Leptogium* (sects. *Leptogium*, *Leptogiopsis* and *Homodium*) in India. *Geophytology* 8(2): 189–204.

- Bungartz F. 2008. Cyanolichens of the Galapagos Islands – the genera *Collema* and *Leptogium*. *Sauteria* 15: 139–158.
- Cao J, Liu HJ, Deng H. 2012. A hairy species of *Leptogium* new to Mainland China. *Journal of Fungal Research* 10(4): 213–215.
- Jaklitsch WM, Baral H-O, Lücking R, Lumbsch HT. 2016. Syllabus of plant families - Adolf Engler's Syllabus der Pflanzenfamilien, Part 1/2, *Ascomycota*. Gebrüder Borntraeger Verlagsbuchhandlung, Berlin. 322 p.
- Jatta A. 1902. Licheni Cinesi raccolti allo Shen-si negli anni 1894-1898 dal. rev. Padre Missionario G. Giraldi. *Nuovo Giornale Botanico Italiano*, ser. 2, 9: 460-481.
- Jørgensen PM. 1994. Further notes on European taxa of the lichen genus *Leptogium*, with emphasis on the small species. *Lichenologist* 26(1): 1–29. <https://doi.org/10.1006/lich.1994.1001>
- Jørgensen PM. 1997. Further notes on hairy *Leptogium* species. *Symbolae Botanicae Upsalienses* 32(1): 113–130.
- Jørgensen PM, Nash TH III. 2004. *Leptogium*. 330–350, in: TH Nash III & al. (eds). *Lichen flora of the Greater Sonoran Desert Region*. Vol. 2. Lichens Unlimited, Arizona State University, Tempe, Arizona.
- Kashiwadani H, Inoue M, Moon KH, Sasaki K. 2002. Lichens of Ozegahara Moor and its adjacent areas, central Japan. *Memoirs of the National Science Museum* 38: 71–93.
- Kirk PM, Cannon PF, Minter DW, Stalpers JA (eds). 2008. *Ainsworth & Bisby's dictionary of the fungi*. 10th edition. CABI International, Wallingford Oxon. 771 p. <https://doi.org/10.1079/9780851998268.0000>
- Liu HJ, Guan S. 2012. A new hairy species of *Leptogium* (*Collemataceae*) from China. *Mycotaxon* 119: 413–417. <https://doi.org/10.5248/119.413>
- Liu HJ, Wu QF, Chen Z. 2010. The lichen order *Peltigerales* from Mt. Liupan, Ningxia, Northwest China. *Journal of Fungal Research* 8(4): 194–199.
- Liu HJ, Cao J, Guan S, Wu QF. 2012. Three non-hairy species of *Leptogium* from China. *Mycotaxon* 122: 483–490. <https://doi.org/10.5248/122.483>
- Liu HJ, Cao J, Guan S, Wu QF. 2013. A non-hairy species of *Leptogium* new to China. *Chenia* 11: 39–42.
- Liu XD, Deng H, Liu HJ. 2014. Two new records of *Leptogium* from China. *Journal of Fungal Research* 12(4): 206–209.
- Liu HJ, Xi MQ, Hu JS, Wu QF. 2015. Two new species and a new record of *Leptogium* from China. *Mycotaxon* 130(2): 471–478. <https://doi.org/10.5248/130.471>
- Lumbsch HT, Huhndorf SM. 2007. Outline of *Ascomycota* – 2007. *Myconet* 13: 1–58.
- Otálora MAG, Jørgensen PM, Wedin M. 2014. A revised generic classification of the jelly lichens, *Collemataceae*. *Fungal Diversity* 64(1): 275–293. <https://doi.org/10.1007/s13225-013-0266-1>
- Sierk HA. 1964. The genus *Leptogium* in North America, north of Mexico. *Bryologist* 67(3): 245–317. <https://doi.org/10.2307/3240893>
- Wang HY, Ren Q, Li HM, Wang HY, Zhao ZT. 2010. Five lichens of *Leptogium* new to China. *Mycotaxon* 111: 161–166. <https://doi.org/10.5248/111.161>
- Wei JC. 1991. *An enumeration of lichens in China*. International Academic Publishers, Beijing. 278 p.
- Xi MQ, Liu HJ. 2014. Two species of *Leptogium* new to Asia. *Journal of Fungal Research* 12(2): 71–74.
- Zahlbruckner A, 1930. Lichenes (Übersicht über sämtliche aus China bekannten Flechten). In: Handdel-Mazzetti H (ed.) *Symbolae sinicae botanisch ergebnisse der expedition der akademie der wissenschaften in wien nach Südwest-China 1914-1918*, Teil 3. Julius Springer Verlag, Wien. 254 p. <https://doi.org/10.1007/978-3-7091-4178-6>