

## SHORT COMMUNICATION

# A new depsidone of *Lobaria pulmonaria* with acetylcholinesterase inhibition activity

Boris Pejin<sup>1,2,3</sup>, Giuseppina Tommonaro<sup>1</sup>, Carmine Iodice<sup>1</sup>, Vele Tesevic<sup>2</sup>, Vlatka Vajs<sup>4</sup>, and Salvatore De Rosa<sup>1</sup>

<sup>1</sup>National Research Council of Italy Institute of Biomolecular Chemistry, CNR-ICB, Pozzuoli (Naples), Italy, <sup>2</sup>Department of Organic Chemistry, Faculty of Chemistry, University of Belgrade, Serbia, <sup>3</sup>Department of Life Sciences, Institute for Multidisciplinary Research, University of Belgrade, Serbia, and <sup>4</sup>Center of Chemistry, Institute of Chemistry, Technology and Metallurgy, University of Belgrade, Serbia

### Abstract

The phytochemical investigation conducted on a foliose lichen, *Lobaria pulmonaria* (L.) Hoffm. (Lobariaceae), led to the isolation of a new depsidone in the form of its diacetate derivative which showed a moderate acetylcholinesterase inhibition activity (1 µg) *in vitro*. This is the first record of identified depsidone structure in searching for these inhibitors.

**Keywords:** Lichen, secondary metabolite, biological activity

## Introduction

Most of the secondary metabolites which are so characteristic of lichens are derived from acetyl-polymalonyl pathway. Indeed, the large majority of depsides, depsidones, dibenzofurans, usnic acids and depsones are peculiar to lichens<sup>1</sup>.

Depsidones have an ether linkage in addition to the ester linkage of the depsides, resulting in a rigid polycyclic system. Therefore, they are based on an 11*H*-dibenzo[*b,e*] [1,4]dioxepin-11-one ring system<sup>2</sup>.

As part of our continuing search on acetylcholinesterase (AChE) inhibitors<sup>3</sup>, a phytochemical investigation was conducted on a foliose lichen, *Lobaria pulmonaria* (L.) Hoffm. (Lobariaceae). The chemical nature of *L. pulmonaria* has been studied by various authors including C. P. Culberson who, when reviewing *Lobaria* species, referred to it as having an interesting distribution of varieties and chemical types containing different combinations of β-oricinol depsidones<sup>4,5</sup>.

## Materials and methods

### Biological material

The lichen *L. pulmonaria* (L.) Hoffm. (Lobariaceae) was collected from *Fagus sylvatica* on the mountain Zelengora

(Bosnia and Herzegovina) in July 2009. Voucher specimen has been deposited in the Herbarium of the Institute of Botany, University of Belgrade, Serbia (BEOU 5997).

### General experimental procedures

<sup>1</sup>H- and <sup>13</sup>C-NMR spectra were recorded at the NMR Service of the Institute for Biomolecular Chemistry of National Council Research of Italy (CNR-ICB) on a Bruker Avance-400 spectrometer operating 400 and 100 MHz, respectively, using an inverse probe fitted with a gradient along the Z-axis, in CDCl<sub>3</sub>, using the solvent signal as an internal standard. Thin-layer chromatography was carried out on precoated silica gel 60 F254 (0.25 mm, Merck, Darmstadt, Germany). LRMS and HRMS were recorded on a JEOL JMS D-300 and an AEI MS-50, respectively.

### Extraction and isolation

Before extraction, the lichen was carefully inspected for contaminants. Air-dried parts of *L. pulmonaria* (70 g) were ground and extracted three times with CHCl<sub>3</sub>, CHCl<sub>3</sub>-MeOH 1:1, MeOH, and MeOH-H<sub>2</sub>O 1:1, respectively, (500 mL each) at room temperature, for up to 1 day each, with the extractives pooled and then evaporated *in vacuo*. The dried CHCl<sub>3</sub>-MeOH (1:1) extract (5.81 g) was dissolved in H<sub>2</sub>O (50 mL) and partitioned

Address for Correspondence: Dr. Boris Pejin, Department of Life Sciences, University of Belgrade, Institute for Multidisciplinary Research, Belgrade, Serbia. E-mail: borispejin@imsi.rs; brspjn@gmail.com

(Received 10 February 2012; revised 07 March 2012; accepted 08 March 2012)

sequentially with  $\text{CHCl}_3$  ( $3 \times 50$  mL) and *n*-BuOH ( $3 \times 50$  mL). The crude insoluble colored residue (0.46 g), obtained after the partition, was classified as fraction rich in epsilons, by means of its spectroscopic data and typical chromatographic profile. In order to further characterize the residue, it was dissolved in 5 mL of pyridine and 1 mL of acetic anhydride (77 mg) and was refluxed for 2 h. After evaporating to dryness, the residue was chromatographed on a silica gel column and eluted with the system of petroleum ether–diethyl ether (gradient, starting from petroleum–ether/diethyl ether 1:1) to yield a mixture of acetylated depsidones (24 mg), which was further separated with thin-layer chromatography using  $\text{CHCl}_3$  to give a new depsidone in the form of its diacetate derivative (3 mg; 0.0043% of dry weight).

A new depsidone (Figure 1): white, amorphous powder; IR (ATR)  $\nu_{\text{max}}$  1781, 1743, 1702, 1606  $\text{cm}^{-1}$ ;  $^1\text{H-NMR}$  (DMSO- $d_6$ , 400 MHz)  $\delta$  10.41 (1H, s, H-9), 7.35 (1H, s, H-6'), 6.79 (1H, s, H-5), 4.00 (3H, s,  $0\text{CH}_3$ -4), 2.58 (3H, s, H-8), 2.43 (3H, s,  $0\text{COCH}_3$ -1'), 2.30 (3H, s, H-8'), 2.23 (3H, s,  $0\text{COCH}_3$ -2');  $^{13}\text{C-NMR}$  (DMSO- $d_6$ , 100 MHz)  $\delta$  186.4 (CHO, C-9), 168.9 (OCO, C-2'), 168.4 (OCO, C-1'), 164.6 (C, C-4), 163.9 (COO, C-7), 160.2 (C, C-2), 151.7 (C, C-6), 148.9 (C, C-2'), 144.7 (C, C-1'), 142.7 (C, C-4'), 134.1 (C, C-5'), 90.8 (C, C-6'), 116.2 (C, C-3'), 114.8 (C, C-1), 114.4 (CH, C-3), 112.2 (CH, C-5), 56.7 (C,  $0\text{CH}_3$ -4), 22.4 ( $\text{CH}_3$ , C-8), 20.7 ( $0\text{COCH}_3$ , C-2'), 20.4 ( $0\text{COCH}_3$ , C-2'), 10.3 ( $\text{CH}_3$ , C-8'); ESIMS  $m/z$  416  $[\text{M} + \text{H}]^+$  (calcd for  $\text{C}_{21}\text{H}_{19}\text{O}_9$  415,1029).

### Biological assay

Acetylcholinesterase inhibition was performed dissolving the sample in MeOH at a concentration of 1 mg/mL. From this main solution was performed a serial dilution in order to obtain lower concentration of the sample (0.1; 0.01; 0.001 mg/mL), and 10  $\mu\text{L}$  of each solution was applied to TLC plates to test 10, 1, 0.1, and 0.01  $\mu\text{g}$  of the sample to detect the minimum concentration that inhibited AChE. Galanthamine was used as positive control. The assay was carried out as described<sup>6</sup>. It is a simple and rapid bioautographic enzyme assay. The test relies on the cleavage by AChE of 1-naphthyl acetate to form 1-naphthol, which in turn reacts with

Fast Blue B salt to give a purple-coloured diazonium dye. Briefly, a stock solution of AChE (1000 U in 150 mL of Tris-hydrochloric acid buffer pH 7.8) was obtained, which was stabilized adding bovine serum albumin (150 mg). A 10  $\mu\text{L}$  aliquot of each solution of the sample was applied to the TLC plates, dried to remove the solvent, and then sprayed with enzyme stock solution. For incubation of the enzyme, the plate was kept at 37°C for 20 min in a humid atmosphere. For the detection of the enzyme, solutions of 1-naphthyl acetate (250 mg in 100 mL of EtOH) and of Fast Blue B salt (400 mg in 160 mL of distilled  $\text{H}_2\text{O}$ ) were mixed and sprayed onto the plate. Acetylcholinesterase inhibition activity was detected by a white spot on a purple background after 1–2 min.

### Results and discussion

High-resolution mass spectrometry established the molecular formula of a new depsidone as  $\text{C}_{21}\text{H}_{18}\text{O}_9$ . Its structure followed from 1-D and 2-D NMR spectra. The  $^1\text{H-NMR}$  spectrum showed five three-proton singlets between  $\delta$  2.0 and 4.0, two one-proton singlets between  $\delta$  6.7 and 7.4, and an aldehyde proton at  $\delta$  10.41; two C-methyl resonances ( $\delta$  2.23 and 2.43) were specific signals of the identified depsidone molecule. The  $^{13}\text{C-NMR}$  and DEPT spectra exhibited 21 carbon signals. A combination of 2-D NMR experiments (COSY, NOESY, HSQC and HMBC) allowed us to assign all signals in the  $^1\text{H}$ - and  $^{13}\text{C}$ -NMR spectra.

The AChE inhibition test<sup>6,7</sup> showed a moderate activity (1  $\mu\text{g}$ ) for a new depsidone in the form of its diacetate derivative. In comparison, the alkaloid galanthamine<sup>8</sup> inhibited the enzyme at 0.01  $\mu\text{g}$ , while the same mixture of depsidones in its acetylated and methylated forms were also active, 0.5 and 2  $\mu\text{g}$ , respectively<sup>9,10</sup>. On the other hand, the mixture in its natural nonmodified form as well as isolated stictic acid (common depsidone for *Lobaria* sp.) did not show any activity in the AChE test on TLC plate.

### Acknowledgements

The assistance of Mr. V. Mirra of NMR Service CNR-ICB is gratefully acknowledged.

### Declaration of interest

This work was supported by the Ministry of Science and Technological Development of the Republic of Serbia (Research grant Nos. 172053 and 173040).

### References

1. Thomas H. Nash III. (2008). *Lichen Biology*. Cambridge University Press.
2. Neamati N, Hong H, Mazumder A, Wang S, Sunder S, Nicklaus MC et al. Depsides and depsidones as inhibitors of HIV-1 integrase: Discovery of novel inhibitors through 3D database searching. *J Med Chem* 1997;40:942–951.




Figure 1. A new depsidone in the form of its diacetate derivative.

3. Pejin B, Iodice C, Tommonaro G, De Rosa S. Synthesis and biological activities of thio-avarol derivatives. *J Nat Prod* 2008;71:1850-1853.
4. González AG, Bermejo Barrera J, Rodríguez Pérez EM, Hernández Padrón CE. Depsidones from *Lobaria pulmonaria* and their chemotaxonomic importance. *Biochem System Ecol* 1994;22:583-586.
5. Culberson CF. Chemical studies in the genus *Lobaria* and the occurrence of a new tridepside, 4-*O*-methylgyrophoric acid. *The Bryologist* 1969;72:19-27.
6. Marston A, Kissling J, Hostettmann K. A rapid TLC bioautographic method for the detection of acetylcholinesterase and butyrylcholinesterase inhibitors in plants. *Phytochem Anal* 2002;13:51-54.
7. Rhee IK, van de Meent M, Ingkaninan K, Verpoorte R. Screening for acetylcholinesterase inhibitors from Amaryllidaceae using silica gel thin-layer chromatography in combination with bioactivity staining. *J Chromatogr A* 2001;915:217-223.
8. Scott LJ, Goa KL. Galantamine: A review of its use in Alzheimer's disease. *Drugs* 2000;60:1095-1122.
9. Pejin B, Tommonaro G, Iodice C, Tesevic V, Vajs V. Acetylcholinesterase inhibition activity of acetylated depsidones from *Lobaria pulmonaria*. *Nat Prod Res* (In Press). DOI: 10.1080/14786419.2011.585989.
10. Pejin B, Tommonaro G, Iodice C, Tesevic V, Vajs V. Acetylcholinesterase inhibition activity of a mixture of methylated depsidones from the lichen *Lobaria pulmonaria*. *Asian J Chem* 2012;24:3261-3262.