

TORREYA

A MONTHLY JOURNAL OF BOTANICAL NOTES AND NEWS



JOHN TORREY, 1796-1873

LIBRARY
NEW YORK
BOTANICAL
GARDEN.

EDITED FOR

THE TORREY BOTANICAL CLUB

BY

NORMAN TAYLOR

Volume XIII

NEW YORK

1913

Found by Mr. George N. Rohwer at station No. 14, Miocene shales of Florissant, Colorado. This may not be a *Phalaris*, but it is at least suggestive of that genus. The specific name is a fanciful one, given in allusion to the parallel straight lines.

SHORTER NOTES

THE GENUS MALPIGHIA IN JAMAICA.—Heretofore we have known five species of *Malpighia* from Jamaica, West Indies, together with the apparently trustworthy report of a sixth one, namely *Malpighia incana* Mill. The species positively known to occur on the island, and also recently collected there, are *Malpighia glabra* L., *M. puniceifolia* L., *M. fucata* Ker., *M. urens* L., and *M. biflora* Poir. Jamaica is the type locality for both *Malpighia glabra* L. and *M. urens* L. Recently specimens of an additional species have come to hand. It may be described as follows:

Malpighia Harrisii Small, sp. nov. A slender shrub commonly about 3.5 m. tall, with sparingly pubescent twigs: leaves glabrous, at least ultimately so, above the petiole; blades oval, oblong, or ovate, 6–12 cm. long, obtuse to acute, or individually notched at the apex, entire, bright-green above, paler-green beneath, rounded at the base, short (5–8 mm.) -petioled: cymes sessile, many-flowered, the branches sparingly pubescent, at least when young: calyx with 6 glands; sepals oblong or ovate-oblong, about 3 mm. long, obtuse, twice as long as the glands or less: corolla rose-color, about 2.5 cm. wide, the larger petals 12–13 mm. long: androecium with 2 very large filaments: gynoecium with unequal styles, the 2 posterior ones much stouter and longer than the anterior one: drupes not seen.

In Peckham Woods, Upper Clarendon, Jamaica, collected by William Harris, September 27, 1912, III 89.

This species appears to be most closely related to *Malpighia urens*. It differs from the latter species in the glabrous, larger, broader, and longer-petioled leaf-blades, and in the larger corollas.

J. K. SMALL

A FURTHER NOTE ON THE LINNEAN HERBARIUM. Dr. B. Daydon Jackson writes me (Jan. 2, 1913) that No. "57 *Cocci-*

ferus" is represented in the Linnean Herbarium, in which I failed to find it (Bull. Torr. Bot. Club 39: 201. 1912). In his recent Index to the Linnean Herbarium (Suppl. Proc. Soc. 96. 1912) he records the species *Lichen cocciferus* L. as represented by a type specimen.

R. HEBER HOWE, JR.

THOREAU MUSEUM,
CONCORD, MASS.

A PLANT NEW TO THE STATE OF NEW YORK AND THE LOCAL FLORA RANGE.—It is here worthy of record that Miss F. A. Mulford collected in July, 1903, at Arkville, Delaware Co., N. Y., specimens of the musk-root *Adoxa Moschatellina* L. This is the first record of this plant's occurrence either in New York State or the local flora range. The nearest previous stations for this species are in Arctic America and Iowa. The discovery of this rare plant at an elevation of about 1,400 ft. in the Catskills is a noteworthy addition to the list of local flora plants. Specimens of this plant from Arkville are in the herbarium of the Brooklyn Botanic Garden and in Miss Mulford's herbarium.

NORMAN TAYLOR

AN INVESTIGATION OF "LACINIARIA SCARIOSA."—As already known at several botanical institutions, the writer has been engaged for many years, as time permitted, in the elaboration of the laciniarias of the *scariosa* group. A large amount of material has been assembled at the National Herbarium by acquisition and loan, the writer himself having collected over six hundred specimens. In all, not less than seventeen hundred specimens have been examined. This material represents beyond all controversy an immense number of species and these, upon due comparison, are seen to fall into natural classes.

It will easily be understood that no superficial study based on local and scattering specimens can secure a scientific disposition of these plants, and the disjointed publication of names, the reference of which can only be known by consulting the type material, adds complication to an already sufficiently complicated matter. I therefore bespeak the patience of the botanical public