

Subnulatones A and B, new *trans*-decalin polyketides from the cultured lichen mycobionts of *Pseudopyrenula subnudata*

Thuc-Huy Duong^{a,b}, Hung-Huy Nguyen^c, Thanh-Trung Le^d, Thanh-Nha Tran^e, Jirapast Sichaem^{f,*}, Thanh-Trung Nguyen^g, Thi-Phuong Nguyen^h, Dinh-Tri Mai^{i,j}, Huu-Hung Nguyen^k, Hoang-Duy Le^{l,*}

^a Department for Management of Science and Technology Development, Ton Duc Thang University, Ho Chi Minh City, Viet Nam

^b Faculty of Applied Sciences, Ton Duc Thang University, Ho Chi Minh City, Viet Nam

^c Department of Inorganic Chemistry, University of Science, Ha Noi National University, 19 Le Thanh Tong Street, Hoan Kiem District, Ha Noi City, Viet Nam

^d Hue University of Science, Hue City, Viet Nam

^e Faculty of Chemistry, University of Education, 280 An Duong Vuong Street, District 5, Ho Chi Minh City, Viet Nam

^f Faculty of Science and Technology, Thammasat University Lampang Campus, Lampang 52190, Thailand

^g Institute of Research and Development, Duy Tan University, Da Nang 550000, Viet Nam

^h Faculty of Biotechnology, Nguyen Tat Thanh University, 300A Nguyen Tat Thanh, District 4, Ho Chi Minh City, Viet Nam

ⁱ Graduate University of Science and Technology, Vietnam Academy of Science and Technology, 18 Hoang Quoc Viet, Cau Giay, Ha Noi, Viet Nam

^j Institute of Chemical Technology, Vietnam Academy of Science and Technology, 01 Mac Dinh Chi, Ho Chi Minh City, Viet Nam

^k Faculty of Environment and Biotechnology, Van Lang University, 45 Nguyen Khac Nhu, District 1, Ho Chi Minh City, Viet Nam

^l Pham Van Dong University, Quang Ngai Province, Viet Nam

ARTICLE INFO

Keywords:

Pseudopyrenula subnudata
Subnulatones A and B
trans-decalin polyketide
Cytotoxicity

ABSTRACT

Chemical investigation of the cultured polyspore-derived mycobionts of a *Pseudopyrenula subnudata* lichen led to the isolation of two new compounds, subnulatones A and B (**1** and **2**), together with four known compounds, 1-(2-hydroxy-1,2,6-trimethyl-1,2,4a,5,6,7,8,8a-octahydronaphthalen-1-yl)ethanone (**3**), libertalide C (**4**), aspermytin A (**5**), and 6,7-dimethoxy-4-hydroxymellin (**6**). Their chemical structures were elucidated by extensive 1D and 2D NMR analysis and high resolution mass spectroscopy, and comparisons were made with the literature. The absolute configuration of **1** was defined unambiguously using single crystal X-ray crystallography. Compound **1** represents the first dimeric decalin polyketide to be found in nature. The *in vitro* cytotoxicity of **1** against two cancer cell lines (K562 and MCF-7) was evaluated. Compound **1** showed moderate cytotoxic activity with IC₅₀ values of 23.5 ± 1.0 and 51.9 ± 1.4 μM, respectively.

1. Introduction

Lichens are symbiotic associations of an algal or cyanobacterial photobiont and a fungal mycobiont, and produce a range of bioactive metabolites. However, the metabolites that are extracted from cultured lichen mycobionts are not detectable in natural lichens under stressed conditions. Phytochemical investigation of mycobionts cultured from Vietnamese lichens identified bioactive compounds with unique scaffolds [1–3]. In our search for metabolites of crustose lichen mycobionts, the mycobionts of *Pseudopyrenula subnudata* lichen were isolated and cultivated on MY10 at 18°C in the dark over several months. The colonies were then harvested and extracted with EtOAc. The extract was separated by a combination of chromatographic procedures to afford two new compounds, subnulatones A and B (**1** and **2**), along with four

known compounds, 1-(2-hydroxy-1,2,6-trimethyl-1,2,4a,5,6,7,8,8a-octahydronaphthalen-1-yl)ethanone (**3**) [4], libertalide C (**4**) [5], aspermytin A (**5**) [6], and 6,7-dimethoxy-4-hydroxymellin (**6**) [7]. The structures of these compounds were determined by extensive spectroscopic analysis (1D, 2D-NMR, and HRESIMS) and single-crystal X-ray crystallography. Compound **1** was assessed for its *in vitro* cytotoxicity against the MCF-7 (breast cancer) and K562 (chronic myelogenous leukemia) cell lines.

2. Experimental

2.1. General experimental procedures

NMR spectra were measured on Bruker Avance II (500 MHz for ¹H

* Corresponding authors.

E-mail addresses: duongthuchuy@tdtu.edu.vn (T.-H. Duong), Jirapast@tu.ac.th (J. Sichaem), lhduy@pdu.edu.vn (H.-D. Le).

<https://doi.org/10.1016/j.fitote.2020.104512>

Received 20 December 2019; Received in revised form 12 February 2020; Accepted 12 February 2020

Available online 13 February 2020

0367-326X/© 2020 Elsevier B.V. All rights reserved.

NMR and 125 MHz for ^{13}C NMR) and Varian Mercury-400 Plus NMR (400 MHz for ^1H NMR and 100 MHz for ^{13}C NMR) spectrometers with TMS as internal standard. Chemical shifts were expressed in ppm with reference to the residual protonated solvent signals (chloroform- d_1 with δ_{H} 7.26 and δ_{C} 77.16, and dimethylsulfoxide- d_6 with δ_{H} 2.50 and δ_{C} 39.52). Crystallographic analyses of suitable single crystals were performed using an automatic diffractometer (Bruker Model D8 Quest) with graphite-monochromated Mo-K α radiation ($\lambda = 0.71073 \text{ \AA}$) at 100 K. The HRESIMS were recorded on a HRESIMS Bruker MicroTOF. TLC was carried out on precoated silica gel 60 F $_{254}$ or Sephadex LH-20 and spots were visualized using UV $_{254\text{nm}}$ and UV $_{365\text{nm}}$ lamps. Gravity column chromatography was performed with silica gel 60 (0.040–0.063 mm).

2.2. Fungal isolation, cultivation, and identification

Specimens of *P. subnudata* Müll. Arg. were collected from tree bark in Quang Ngai City (15.10437 N; 108.80656 E), Vietnam (approx. 9 m alt.), in March 2017 by D.H. Le. The voucher specimens were identified by Dr. Vo Thi Phi Giao (University of Sciences, Ho Chi Minh City, Vietnam) and deposited at the University of Sciences, Ho Chi Minh City, Vietnam (registration No. QNg170301). Mycobionts were obtained from the spores discharged from the apothecia of a thallus, and were cultivated in test tubes containing modified MY10 medium (malt extract 10 g, yeast extract 4 g, sucrose 100 g, agar 15 g, H $_2$ O 1 L, pH 7) at 18 °C in the dark. After cultivation for 4–8 months, the colonies were harvested.

2.3. Extraction and isolation

The harvested colonies (dry weight 31.5 g) were extracted with EtOAc (3 \times 100 mL each) at room temperature, and the combined extracts were concentrated under reduced pressure to give a 608.8 mg residue. The EtOAc extracts were subjected to silica gel column chromatography (CC) and eluted by the solvent system *n*-hexane-EtOAc with increasing ethyl acetate ratios to yield eighteen fractions: EA1 to EA18. Fraction EA3 (12 mg) was subjected to reversed-phase CC using solvent system of MeOH-H $_2$ O (10:1, v/v) as a mobile phase to afford 2 (2 mg), 3 (5.3 mg), and 4 (1.8 mg). Fraction EA9 (173 mg) was used for further isolation through reversed-phase CC using a solvent system of MeOH-H $_2$ O (10:1, v/v), yielding 1 (44 mg). This was crystallized in MeOH, to obtain a single purified crystal. Fraction EA14 (109 mg) was subjected to a silica gel CC eluted with *n*-hexane-CHCl $_3$ -EtOAc-acetone-AcOH (100:20:8:5:2) to give five subfractions (EA14.1-EA14.5). Fraction EA14.3 (25 mg) was selected for reversed-phase CC using MeOH-H $_2$ O (5:1). Three subfractions (EA14.3.1-EA14.3.3) were obtained. Compound 5 was isolated from subfraction EA14.3.1 by crystallization in MeOH. Finally, subfraction EA14.3.3 (5 mg) was purified by a Sephadex LH-20 with MeOH-CH $_2$ Cl $_2$ (1:1, v/v) as an eluent to afford 6 (1.2 mg).

2.3.1. Subnudatone A (1)

Colorless crystal; $[\alpha]_{\text{D}}^{22} + 115$ (c 0.1, MeOH); UV (EtOH) λ_{max} (log ϵ) 204 (2.9), 285 (1.9) nm; ^1H (CDCl $_3$, 500 MHz) and ^{13}C (CDCl $_3$, 125 MHz) NMR data, see Table 1; HRESIMS m/z 493.3321 [M + H] $^+$ (calcd for C $_{32}$ H $_{45}$ O $_4$, 493.33129); 515.3015 [M + Na] $^+$ (calcd for C $_{32}$ H $_{44}$ O $_4$ Na, 515.3137).

2.3.1.1. X-ray crystallographic analysis of 1. Single crystals of 1 were obtained by vapor diffusion of MeOH, in the form of colorless crystals. Crystal analysis was conducted using a D8 Quest Diffractometer with Mo-K α radiation ($\lambda = 0.71073 \text{ \AA}$) at 100.0 [2] K. The structures were solved by the intrinsic phasing method (ShelXT) and refined using a full-matrix least-squares difference Fourier technique (ShelXL). The positions of hydrogen atoms were determined using ‘riding’ mode. The crystallographic data for Compound 1 has been deposited with the

Cambridge Crystallographic Data Centre. Copies of the data can be obtained, free of charge, on application to the Director, CCDC, 12 Union Road, Cambridge CB2 1EZ, UK (fax: 44-(0)1223-336,033, or e-mail: deposit@ccdc.cam.ac.uk). Crystal data of 1: C $_{32}$ H $_{44}$ O $_4$, Mr. = 492.67, orthorhombic, $a = 6.238 [1] \text{ \AA}$, $b = 12.317 [1] \text{ \AA}$, $c = 36.844 [2] \text{ \AA}$, $\alpha = 90.00$, $\beta = 90.00$, $\gamma = 90.00$, $V = 2830.6 [3] \text{ \AA}^3$, space group P2 $_1$ 2 $_1$ 2 $_1$, $Z = 4$, $D_{\text{calcd}} = 1.156 \text{ g/cm}^3$, $\mu = 0.074 \text{ mm}^{-1}$, and $F(000) = 1072.0$. Crystal dimensions: 0.25 \times 0.25 \times 0.10 mm 3 . Independent reflections: 70333 ($R_{\text{int}} = 0.0561$). The final R1 values were 0.0482, $\omega R2 = 0.0876$ ($I > 2\sigma(I)$). The goodness of fit on F2 was 1.066. Flack parameter: 0.2 [3]. CCDC number: 1954725.

2.3.2. Subnudatone B (2)

Colorless oil; $[\alpha]_{\text{D}}^{22} + 315$ (c 0.1, MeOH); UV (EtOH) λ_{max} (log ϵ) 204 (2.5), 283 (1.7) nm; ^1H (CDCl $_3$, 400 MHz) and ^{13}C NMR (CDCl $_3$, 100 MHz), see Table 1; HRESIMS m/z : [M + Na] $^+$ 269.1514 (calcd for C $_{16}$ H $_{22}$ O $_2$ Na, 269.1517).

2.4. Cytotoxicity assay

The cytotoxicity of 1 was evaluated against the MCF-7 (breast cancer) and K562 (chronic myelogenous leukemia) tumor cell lines, cultured in RPMI 1640 medium and in DMEM medium, respectively. The cultures were supplemented with 10% fetal bovine serum (FBS), 100 IU/mL penicillin, and 100 $\mu\text{g/mL}$ streptomycin, and maintained at 37 °C, 5% CO $_2$, and 95% humidity. Viable cells were counted and inoculated into a 96-well plate at densities of 10 4 cells/100 μL /well for MCF-7 and 10 5 cells/100 μL /well for K562. After 24 h, the cells were treated with the two compounds, using doxorubicin as a positive control. They were diluted in the culture media at concentrations of 100, 50, 25, 12.5, 6.25, 3.125, and 0 $\mu\text{g/mL}$, containing 1, 0.5, 0.25, 0.125, 0.0625, 0.03125, and 0% dimethyl sulfoxide (DMSO), respectively. The DMSO was used as a negative control, and culture medium without cells was used as a blank. All experiments were done in triplicate. The plates were incubated at 5% CO $_2$, 95% humidity, and 37 °C for 72 h. 10 μL of 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT, 5 mg/mL stock solution) was added to each well and incubated at 37 °C in 5% CO $_2$ for 3.5 h. 70 μL of Detergent Reagent (10% SDS) was added to each well and the plate was maintained at 37 °C for 16 h. The optical density of each well was read by a scanning multiwell spectrophotometer (Sunrise) at a wavelength of 595 nm. Cell survival was measured by comparing the percentage absorbance with that of the negative control.

3. Results and discussion

Compound 1 was isolated as a colorless crystal. The molecular formula was established to be C $_{32}$ H $_{44}$ O $_4$ on the basis of the positive-ion mode HRESIMS data with a sodiated ion peak at m/z 515.3015 [M + Na] $^+$ (calcd. 515.3137 for C $_{32}$ H $_{44}$ O $_4$ Na). The ^1H NMR and HSQC spectra showed seven olefinic protons (δ_{H} 7.50, 7.45, 7.02, 5.67, 5.46, 5.37, and 5.37, each 1H), four tertiary methyl protons (δ_{H} 1.42, 1.22, 1.26, and 1.17, each 3H), two secondary methyls (δ_{H} 0.91 and 0.93, each 3H), and high field protons in the range 0.85 to 2.83 ppm. The ^{13}C and DEPT NMR spectra revealed 32 carbons, including two ketone carbons (δ_{C} 204.2 and 197.8), seven olefinic methine carbons (δ_{C} 162.6, 135.3, 133.2, 132.4, 130.6, 128.0, and 123.5), five quaternary carbons (δ_{C} 114.4, 87.8, 74.2, 56.2, and 48.3) of which two (δ_{C} 87.8 and 74.2) were oxygenated, six methyl carbons (δ_{C} 28.2, 23.0, 22.6, 22.5, 13.9, and 12.1), six sp 3 methylene carbons (δ_{C} 41.8, 41.2, 35.7, 35.6, 28.2, and 27.7), and six sp 3 methine carbons (δ_{C} 43.7, 42.7, 41.3, 39.1, 33.7, and 33.6). The chemical features were similar to those of 16-carbon polyketides, having a *trans*-fused decalin ring that is common in marine-fungal and lichen-fungal sources [8–11].

The HMBC correlations identified the first spin system, later referred to as subunit A. The HMBC correlation of protons at H-1 (δ_{H}

Table 1
 ^1H and ^{13}C NMR data for compounds **1** and **2** (CDCl_3).

1^a			2^b					
Position	Subunit A δ_{H} , mult (<i>J</i> , Hz)	δ_{C}	Position	Subunit B δ_{H} , mult (<i>J</i> , Hz)	δ_{C}	Position	δ_{H} , mult (<i>J</i> , Hz)	δ_{C}
1	7.35, 1H, s	162.6	1'	7.50, 1H, d (15.5)	135.3	1	6.98, 1H, d (5.6)	157.5
2		114.4	2'	7.02, 1H, d (15.5)	123.5	2	5.22, 1H, d (6.0)	106.6
3		197.8	3'		204.2	3		200.6
4		48.3	4'		56.2	4		48.5
5	1.64, 1H, td (12.0, 2.0)	42.7	5'	1.83, 1H, m	43.7	5	1.63, 1H, td (12.0, 2.5)	42.7
6	2.82, 1H, dq (12.0, 2.5)	27.7	6'	1.71, 1H, m 1.07, 1H, m	28.2	6	2.82, 1H, dq (12.0, 3.2)	27.6
7	1.77, 1H, m 1.78, 1H, m	35.6	7'	1.79, 1H, m 0.83, 1H, m	35.7	7	1.77, 2H, m	35.7
8	1.50, 1H, m	33.6	8'	1.50, 1H, m	33.7	8	1.49, 1H, m	33.6
9	1.80, 1H, m 0.85, 1H, m	41.8	9'	1.80, 1H, m 0.85, 1H, m	41.2	9	1.80, 1H, m 0.85, 1H, m	41.2
10	1.93, 1H, m	41.3	10'	1.87, 1H, m	39.1	10	1.92, 1H, m	41.1
11	5.46, 1H, dd (10.0, 2.5)	132.4	11'	5.37, 1H, brs	130.6	11	5.75, 1H, dd (10.0, 2.8)	131.8
12	5.66, 1H, dd (10.0, 2.0)	128.0	12'	5.37, 1H, brs	133.2	12	5.42, 1H, dd (10.0, 2.0)	128.7
13		87.8	13'		74.2	13		86.3
14	1.42, 3H, s	23.0	14'	1.17, 3H, s	28.2	14	1.44, 3H, s	22.6
15	1.22, 3H, s	13.9	15'	1.26, 3H, s	12.1	15	1.28, 3H, s	14.3
16	0.91, 3H, d (6.5)	22.5	16'	0.93, 3H, d (6.5)	22.6	16	0.91, 3H, d (6.8)	22.2

^a ^1H (500 MHz) and ^{13}C (125 MHz) NMR.

^b ^1H (400 MHz) and ^{13}C (100 MHz) NMR.




Fig. 1. Chemical structures of compounds **1–6**.

7.35) to carbon at C-3 (δ_{C} 197.8) and C-2 (δ_{C} 114.4); H_3 -15 (δ_{H} 1.22) to C-3 (δ_{C} 197.8) and C-4 (δ_{C} 48.3); and H_3 -14 (δ_{H} 1.42) to C-4 (δ_{C} 48.3) and C-13 (δ_{C} 87.8) defined the connectivity as C-1-C-2-C-3-C-4-C-13. The downfield ^{13}C chemical shift of C-1, together with the key HMBC correlation of H-1 to C-13, indicated ether linkage between C-1 and C-13. Furthermore, HMBC correlations of H_3 -14 to C-12 (δ_{C} 128.0), and of both olefinic protons H-11 (δ_{H} 5.46) and H-12 (δ_{H} 5.66) to C-13, established the vicinity of these groups. The rest of the subunit A could be defined from ^1H - ^1H COSY correlations of H-11/H-10/H-7/H-9/H-8/ H_3 -16 and H-8/H-7/H-5. Finally, HMBC correlations of H-5 (δ_{H} 1.64) to

C-4 and C-3 validated the C-3-C-4-C-5 connectivity. The large coupling constant of H-5 (12.0 Hz) confirmed the *trans*-decalin conformation of subunit A, further supported by the NOESY correlations. NOESY correlations of H_3 -14 (δ_{H} 1.42)/H-5 (δ_{H} 1.64) and H_3 -15 (δ_{H} 1.22)/H-10 (δ_{H} 1.93) were observed, while the absence of NOESY correlations between the two methyls indicated the *syn*-orientation of H-5/ H_3 -14 and H-10/ H_3 -15. The NMR data on subunit A were strongly reminiscent of those of versiol [8] with the exception of an additional double bond at C-1 and C-2. Accordingly, subunit A was characterized as shown in Fig. 1.




Fig. 2. Key ^1H - ^1H COSY, HMBC, and NOESY correlations of compounds 1 and 2.

In subunit B, two *trans* coupled olefinic protons at δ_{H} 7.50 (H-1') and 7.02 (H-2') gave HMBC correlation to C-2. Conversely, proton H-1 (δ_{H} 7.35) showed HMBC correlation to C-1', indicating connectivity between subunits A and B at C-2. HMBC cross peaks of H-1' (δ_{H} 7.50) and H₃-15' (δ_{H} 1.26) to C-3 and C-4, and H₃-14' (δ_{H} 1.17) to C-4' (δ_{C} 56.2), suggested connectivity through C-1' to C-4'. Moreover, the methyl H₃-14' gave an HMBC cross peak to C-12' (δ_{C} 133.2), indicating a vicinal C-11' and C-12' double bond. COSY correlations from H-5' to H-12' validated the remaining signals of subunit B. In this subunit, NOESY correlations of H-5'/H-1'/H₃-14' defined their co-facial side. Likewise, a NOESY cross peak between methyl H₃-15' and H-5' indicated that they were in the same orientation. These chemical features were very similar to those of aspermytin A (5), with the exception of an additional double bond at C-1 and C-2.

The absolute stereochemistry of 1 was determined using single-crystal X-ray crystallography. A qualified single crystal was successfully obtained and an ORTEP diagram identified a *trans*-decalin conformation in both subunits A and B as well as other chiral centers (see Fig. 3). The chemical structure of 1 was elucidated as that of subnudatone A, and is shown as Fig. 1.

Compound 2 was obtained as colorless oil. Its molecular formula was determined as C₁₆H₂₂O₂ by the positive HRESIMS measurement through the ion peak at m/z 269.1514 (calcd. 269.1517 for C₁₆H₂₂O₂Na). The ^1H NMR and HSQC spectra showed two singlet methyl groups [δ_{H} 1.44 (3H, s, H₃-14) and 1.28 (3H, s, H₃-15)], four olefinic protons [δ_{H} 6.98 (1H, d, J = 5.6 Hz, H-1), 5.65 (1H, dd, J = 10.0, 2.8 Hz, H-12), 5.42 (1H, dd, J = 10.0, 2.0 Hz, H-11), 5.22 (1H, d, J = 6.0 Hz, H-2), one secondary methyl group [δ_{H} 0.88 (3H, d, J = 6.8 Hz, H₃-16)], and protons in the range 1.00 to 3.11 ppm. The ^{13}C and DEPT NMR spectra revealed 16 carbon resonances attributable to one ketone carbonyl group (δ_{C} 200.6), three methyl carbons (δ_{C} 22.8, 14.5, and 14.3), three methylene carbons (δ_{C} 41.4, 35.7, and 27.6), four sp² methine carbons (δ_{C} 157.5, 131.8, 128.7, and 106.6), three sp³ methine carbons (δ_{C} 42.7, 41.2, and 33.6), and two quaternary carbons (δ_{C} 86.3 and 48.5) of which one (δ_{C} 86.3) was oxygenated. NMR data of 2 were highly similar to those of the subunit A of 1. Indeed, the HMBC correlations of protons at H-1 (δ_{H} 6.98) to carbon at C-3 (δ_{C} 200.6), C-2 (δ_{C} 106.6); H₃-15 (δ_{H} 1.28) to C-3 and C-4 (δ_{C} 48.5); H₃-14 (δ_{H} 1.44) to C-4 (δ_{C} 48.5) and C-13 (δ_{C} 86.3) defined the connectivity through C-1-C-2-C-3-C-4-C-13. The relative configuration of 2 was defined due to the analysis of the coupling constant values and NOESY correlations. Particularly, the *trans*-decalin conformation of 2 was determined by the large coupling constant of H-5 (12.0 Hz) and further backed up by NOESY data (see Fig. S14). Accordingly, the structure of 2 was established and named as subnudatone B.

To date, over 40 *trans*-decalin polyketides with similar scaffolds [6,8–11] have been isolated. Very rare decalins having a C-1-C-2 double

bond have been reported, but no dimeric compounds were found. The linkage between monomers A and B made our skeleton unique. To the best of our knowledge, subnudatone A (1) is the first example of a dimer comprising two *trans*-fused decalin moieties. Subnudatone A (1) bears striking similarities to co-isolated compounds, subnudatone B (2) and aspermytin A (5), proposing that 1 should be a C-acylated product of two monomers. First, compound 5 was dehydrated to form the monomer B. Then, this activated monomer enables the cross-coupling reaction with 2 (subunit A) to provide 1 [12]. Compound 3 was synthesized previously by Inoue and coworkers when these authors conducted the total synthesis of aspermytin A [4]. Therefore, this compound was found as a new natural compound.

The cytotoxicity against K562 and MCF-7 cell lines of 1 was investigated. Activity against these two cancer cell lines was moderate, with IC₅₀ values of 23.5 ± 1.0 and 51.9 ± 1.4 μM, respectively.

4. Conclusions

In this study, six compounds (1–6) were isolated from the cultured lichen mycobionts of *P. subnudata* and their structures were elucidated as subnudatones A and B (1 and 2), 1-(2-hydroxy-1,2,6-trimethyl-1,2,4a,5,6,7,8,8a-octahydronaphthalen-1-yl)ethanone (3), libertalide C (4), aspermytin A (5), and 6,7-dimethoxy-4-hydroxymellin (6). To the best of our knowledge, compound 2 is a new compound and represents the first example of *trans*-decalin polyketide with a C-1-C-2 double bond, while compound 1 is the first example of a novel dimer comprising two *trans*-fused decalin skeletons. On the other hand, compound 3 is a new natural compound. Compound 1 showed moderate cytotoxic activity against K562 (IC₅₀ 23.5 ± 1.0 μM) and MCF-7 (IC₅₀ 51.9 ± 1.4 μM) cell lines Fig. 2.




Fig. 3. ORTEP drawing of compound 1.

Declaration of Competing Interest

No potential conflict of interest was declared by the authors.

Acknowledgements

This research was partly supported by International Foundation for Science (IFS) for D. H. Le under grant number I-1-F-6042-1. Thanks are also for Dr. T. P. G. Vo (University of Science, Ho Chi Minh City, Viet Nam) due to identification of the voucher specimens.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.fitote.2020.104512>.

References

- [1] D.H. Le, Y. Takenaka, N. Hamada, H. Miyawaki, T. Tanahashi, A 14-membered macrolide and isocoumarin derivatives from the cultured lichen mycobionts of *Graphis vestitoides*, *Chem. Pharm. Bull.* 61 (2013) 358–362.
- [2] D.H. Le, Y. Takenaka, N. Hamada, Y. Mizushima, T. Tanahashi, Polyketides from the cultured lichen mycobiont of a Vietnamese *Pyrenula* sp, *J. Nat. Prod.* 77 (2014) 1404–1412.
- [3] D.H. Le, Y. Takenaka, N. Hamada, T. Tanahashi Eremophilane-type sesquiterpenes from cultured lichen mycobionts of *Sarcographa tricosia*, *Phytochemistry* 91 (2013) 242–248.
- [4] A. Inoue, M. Kanematsu, S. Mori, M. Yoshida, K. Shishido, An efficient access to aspermytin A and oblongolide C through an intramolecular nitrile oxide–alkene [3+ 2] cycloaddition, *Synlett.* 24 (2013) 61–64.
- [5] C.L. Sun, H. Yan, X.H. Li, X.F. Zheng, H.Y. Liu, Terpenoids from *Chloranthus elatior*, *Nat. Prod. Bioprospect.* 2 (2012) 156–159.
- [6] S. Tsukamoto, S. Miura, Y. Yamashita, T. Ohta, Aspermytin A: a new neurotrophic polyketide isolated from a marine-derived fungus of the genus *Aspergillus*, *Bioorg. Med. Chem. Lett.* 14 (2004) 417–420.
- [7] M.H. Haroon, S.R. Premaratne, M.I. Choudhry, H.R. Dharmaratne, A new β -glucuronidase inhibiting butyrolactone from the marine endophytic fungus *Aspergillus terreus*, *Nat. Prod. Res.* 27 (2013) 1060–1066.
- [8] Y. Fujii, M. Asahara, M. Ichinoe, H. Nakajima, Fungal melanin inhibitor and related compounds from *Penicillium decumbens*, *Phytochemistry* 60 (2002) 703–708.
- [9] H. Guo, T. Feng, Z.H. Li, J.K. Liu, Five new polyketides from the basidiomycete *Craterellus odoratus*, *Nat. Prod. Bioprospect.* 2 (2012) 170–173.
- [10] Y. Ma, J. Li, M. Huang, L. Liu, J. Wang, Y. Lin, Six new polyketide decalin compounds from mangrove endophytic fungus *Penicillium aurantiogriseum* 328, *Mar. drugs* 13 (2015) 6306–6318.
- [11] K. Wijesekera, C. Mahidol, S. Ruchirawat, P. Kittakoop, Metabolite diversification by cultivation of the endophytic fungus *Dothideomycete* sp. in halogen containing media: Cultivation of terrestrial fungus in seawater, *Bioorg. Med. Chem.* 25 (2017) 2868–2877.
- [12] Z.K. Wen, Y.H. Xu, T.P. Loh, Palladium (ii)-catalyzed cross-coupling of simple alkenes with acrylates: a direct approach to 1, 3-dienes through C–H activation, *Chem. Sci.* 4 (2013) 4520–4524.