

**ASSOCIATIONS BETWEEN TREE SNAILS AND CORTICOLOUS LICHENS IN A SECONDARY FOREST IN EASTERN CUBA**Bernardo REYES-TUR<sup>1</sup>✉, Adrián GONZÁLEZ-GUILLÉN<sup>2</sup>, Dania ROSABAL<sup>3</sup> y Alejandro CAPOTE-DANET<sup>1</sup>

1. Departamento de Biología y Geografía, Facultad de Ciencias Naturales y Exactas, Universidad de Oriente, Ave. Patricio Lumumba, Santiago de Cuba 90500, Cuba.

2. Sociedad Cubana de Zoología

3. Centro de Desove del Camarón, UEB Manzanillo, Granma, Cuba.

**ABSTRACT:** Snail-lichen interactions have received little attention in tropical ecosystems. Here, we studied the species richness of corticolous lichens with snail grazing traces on host trees inhabited by the Cuban tree snails, *Polymita venusta*, *Liguus fasciatus* and *Hemistrochus lucipeta* at La Rinconada, eastern Cuba. We sampled lichen specimens from ground level to 2 m, on the trunks of 15 host trees of eight species, in a 6 000 m<sup>2</sup> forest patch, in April 2012. In addition, we sampled snail feces from the three snail species. Most of the tree snail and lichen species were associated with a tree species usually with smooth bark, *Senna atomaria*. We found 30 lichen species, of which 19 were identified to species, three to genus, and eight remain undetermined. Non-metric multidimensional scaling and  $\chi^2$  tests showed that the three snail species had different associations with lichen species. *Liguus fasciatus* is associated with *Physcia sorediosa* and *P. aipolia*; *H. lucipeta* is associated with *Platythecium grammitis* and an unknown lichen, and *P. venusta* is associated with *Buellia spuria* and *Chrysothrix candelaris*. Evidence from snail grazing damage on lichen species, together with ingested spores from six species, suggested that *L. fasciatus* and *P. venusta* are generalist herbivores.

**KEYWORDS:** herbivory, lichen-animal interaction, Gastropoda, Stylommatophora.

**RESUMEN:** ASOCIACIONES ENTRE MOLUSCOS ARBORÍCOLAS Y LÍQUENES CORTÍCOLAS EN UN BOSQUE SECUNDARIO DE CUBA ORIENTAL. En los ecosistemas tropicales las interacciones moluscos-líquenes han recibido escasa atención. En el presente trabajo se estudió la riqueza específica de líquenes cortícolas con huellas de herbivoría de los moluscos arborícolas cubanos,

*Polymita venusta*, *Liguus fasciatus* and *Hemistrochus lucipeta*, en La Rinconada, Cuba oriental. En abril 2012, en un transecto de bosque secundario de 6 000 m<sup>2</sup>, se recolectaron muestras de líquenes hasta 2 m de altura sobre el suelo, en troncos de 15 árboles de 8 especies. Además, se recolectaron heces de las tres especies de moluscos. La mayoría de los moluscos y líquenes utilizaron como substrato árboles de *Senna atomaria*, generalmente con corteza lisa. Se recolectaron 30 especies de líquenes, 19 identificadas hasta especie, 3 hasta género y 8 permanecen sin identificación. El Escalamiento Multidimensional no Paramétrico y las pruebas  $\chi^2$  mostraron que las tres especies de moluscos se asocian de manera diferencial con especies de líquenes. *Liguus fasciatus* estuvo asociado con *Physcia sorediosa* y *P. aipolia*, *H. lucipeta* con *Platythecium grammitis* y una especie indeterminada, mientras *P. venusta* se asoció con *Buellia spuria* y *Chrysothrix candelaris*. La evidencia de las huellas de herbivoría de los moluscos a los líquenes, conjuntamente con la presencia en las heces de esporas de seis especies de líquenes, sugiere que *L. fasciatus* y *P. venusta* son especies herbívoras generalistas.

**PALABRAS CLAVE:** Gastropoda, herbivoría, interacción líquen-animal, Stylommatophora.

**INTRODUCTION**

Snails are key components of terrestrial communities due to their role as detritivores and herbivores (Rollo, 1988; Baur *et al.*, 1992, 1994, 2000; Curry, 1994; Hatzioannou *et al.*, 1994; Cameron, 2016; Price *et al.*, 2016). They are known to exert selective pressures affecting the phenology, morphology and defensive systems of various tree species and their epiphytes, including lichens (Lücking and Bernecker-Lücking, 2000; Speiser, 2001; Asplund and Gauslaa, 2008; Asplund *et al.*, 2010; Boch *et al.*, 2016).

✉ Bernardo REYES-TUR  
breyestur@gmail.com

Publicado en línea el 30 de junio de 2020



Land snails that feed on lichen are often generalists and their snail grazing traces on lichens have been frequently reported (Coker, 1967; Peake and James, 1967; Shachak *et al.*, 1987; Fröberg *et al.*, 1993; Baur *et al.*, 1995, 2000; Lücking and Bernecker-Lücking, 2000; Vatne *et al.*, 2010; Boch *et al.*, 2015; Asplund *et al.*, 2016, 2018). Although there is a growing interest in snail-lichen associations (Lawrey, 1983; Fröberg *et al.*, 2006, 2011; Asplund *et al.*, 2010; Asplund and Wardle, 2013, 2017), most of the focus has been mainly on non-tropical regions despite the remarkable lichen species richness occurring in the tropics (Lücking *et al.*, 2014; Hawksworth and Lücking, 2017). In Cuba, this gap is noteworthy given not only the high diversity of Cuban lichens (Pluntke, 1984; Minter *et al.*, 2001, 2002; Rosabal *et al.*, 2016) but also of Cuban land snails (Hernández-Quinta *et al.*, 2017).

Torre (1950) remarked that the Cuban snails, *Polymita* spp., feed exclusively on lichens and fungi growing epiphytically on their host trees. Díaz-Piferrer (1961) confirmed Torre's approach through an analysis of the stomach content in *Polymita muscarum* (Lea, 1834). This snail species was associated with host trees having a high abundance of epiphyte fungi (Bidart, 1997). Likewise, the most common host tree species of *Polymita venusta* (Gmelin, 1792) also supported the most lichen genera (Reyes-Tur and González-Rodríguez, 2003). *Liguus fasciatus* (Müller, 1774), a tree snail often co-occurring with *Polymita* spp. seems to dis-

play similar feeding preferences for corticolous lichens (*i.e.*, bark-growing), fungi and algae (Pilsbry, 1946; González-Guillén, 2014; González-Guillén *et al.*, 2018). Yet, for the rest of Cuban tree snails, information is scattered, insufficient or missing.

Here, our purpose was to identify the lichen species with grazing traces growing on host trees of the tree snails *P. venusta*, *L. fasciatus* and *Hemitrochus lucipeta* (Poey, 1854) in hills located at La Rinconada, Jiguaní municipality, Granma province, Cuba. We also evaluated the association of the three tree snails with corticolous lichen species, including evidence from snail feces.

## MATERIALS AND METHODS

### STUDY AREA

The studied forest was located at La Rinconada locality, 10 km Southeast of Jiguaní municipality in Granma province, eastern Cuba (Fig. 1). The stand was a secondary forest, as defined by the criteria of Capote and Berazaín (1984). Dominant plant species included *Acacia* sp.; *Bromelia pinguin*, L.; *Comocladia dentata*, Jacq.; *Croton lucidus*, L.; *Cupania* sp.; *Eugenia* sp.; *Lantana* sp.; *Lysiloma* sp.; *Senna atomaria* (L.) Irwin & Barneby, and *Smilax havanensis*, Jacq. The maximum height of these trees and shrubs ranged between 5 and 12 m and they formed a closed canopy with a 50 m<sup>2</sup> total area of harvested-related gaps.




FIGURE 1. Geographical location of study site in Jiguaní municipality, on eastern Cuba.  
 FIGURA 1. Posición geográfica de la localidad de estudio en el municipio Jiguaní, Cuba Oriental.

#### FIELD SAMPLING AND LABORATORY ANALYSIS

Field work was done April 14<sup>th</sup>-15<sup>th</sup>, 2012 in a 6 000 m<sup>2</sup> transect (2 000 m length x 3 m width). We selected 15 living trees of 8 species on which we found evidence for presence of the tree snails *P. venusta*, *L. fasciatus* and *H. lucipeta*. Distance to closest sampled tree was minimum 40 m. Each studied host tree was identified to species. We recorded host tree total height (in m), snail species present, number of snail individuals, and lichen species richness up to 2 m height of the trunk. Each tree was investigated by two researchers for 1 h, the total area sampled was between 1 000 and 2 000 cm<sup>2</sup>.

Corticolous lichens bearing evidence of snail herbivory were sampled for identification in the laboratory. In addition, a sample of snails was captured and housed individually in plastic containers. Their fecal matter was sampled after 20 min and then they were released to their original host trees. Fecal samples were kept in plastic bags at room temperature, between 7-15 days prior to examination with a stereomicroscope in the laboratory. Grazing traces on lichens were ascribed to these gastropods through direct observations and examination of grazed damage by photography of the traces by snail radulae (Fig. 2).

Lichens were identified using specialized keys (Wirth and Hale, 1978; Moberg, 1990; Brodo *et al.*, 2001; Aptroot *et al.*, 2008) and viewing through two Novel microscopes: XSZ-N207 and NTB-2B. Nomenclature followed Mycobank (Crous *et al.*, 2004; Robert *et al.*, 2005, 2013).

#### DATA ANALYSIS

The occurrence frequency of each lichen species was estimated as the percentage of host trees with presence records of each lichen species of the total number of studied trees. A Sørensen similarity matrix was computed from the presence-absence matrix of tree snail and lichen species in sampled host trees. This similarity matrix including both, snails and lichens, was used to conduct a Non-metric multidimensional scaling (NMDS) for the visualization of multivariate patterns. This scaling was accomplished with 25 permutations, establishing as significant interaction criteria the stress values over or equal to 0.1 (Jongman *et al.*, 2001; Quinn and Keough, 2002). To validate the NMDS results, we used the  $\chi^2$  (Chi-square) test to compare the observed frequencies of co-occurrence between the tree snail and lichen species, and the expected frequencies if such co-occurrence was just random. Statistical routines were performed using STATISTICA v. 6.1 (StatSoft, 2003) and PRIMER v. 6.0 (Clarke and Gorley, 2006).

#### RESULTS

##### TREE SNAILS ASSOCIATED WITH HOST TREES

*Liguus fasciatus* was the most frequent snail (n = 38 snail specimens, 73 % of the trees), followed by *P. venusta* (n = 21, 60 %) and *H. lucipeta* (n = 29, 20 %, restricted to *Lysiloma* sp.), with a total of 88 snails over all three species. Over all



FIGURE 2. Different patterns of grazing traces on lichens caused by the radulae of tree snails, A. Large grazing traces and B. Small grazing traces.

FIGURA 2. Diferentes patrones de huellas de herbivoría causadas por las rádulas de moluscos arborícolas a los líquenes, A. Huellas de talla grande y B. Huellas de talla pequeña.

species, abundance of snails per host tree ranged from one to 18 individuals. Grazing traces of tree snails on lichens were recorded for 93 % of the host trees. Most of the trees had smooth bark (53 %) and heights ranged from 2 to 9 m. Tree snails were detected on 8 tree species, with higher frequencies on *Senna atomaria* (40 %) and *Lysiloma* sp. (20 %); these two species commonly had smooth barks and varied in height between 3 and 6 m. Host species with only one occurrence of a tree snail were *Melicoccus bijugatus* Jacq., *Acacia* sp., *Eugenia* sp., *Harrisia eriophora* (Pfeiff.) Britton, *Oxandra laceolata* (Sw.) Baill., and one unidentified tree species without leaves and flowers; however, only the first species had smooth bark.

#### TREE SNAIL-CORTICOLOUS LICHEN ASSOCIATIONS

The corticolous lichen species collected in the studied locality are given in Table 1. There were 30 lichen species belonging to at least 17 genera and 12 families; 19 were identified to species; three to genus, and eight were not determined because they were sterile or lacked spores. Most (56.7 %) were crustose, 40 % foliose and just 3.3 % fruticose. The families with highest species diversity were Graphidaceae (4) and Parmeliaceae (4). Species richness ranged between 1 to 9 lichen species, with 73 % of the trees having from 2 to 6 lichen species. *Senna atomaria* had from 3 to 9 species per tree and hosted 76.7 % of recorded lichen species. Although, most of these lichens were associated with *P. venusta* (83 %,  $n = 30$ ) and *L. fasciatus* (80 %); a lower association was with *H. lucipeta* (17 %).

Higher lichen occurrence frequency in host trees was recorded for *Buellia spuria* (47 %), *Chrysothrix candelaris* (33 %), *Physcia solediosa* (33 %) and *Physcia aipolia* (27 %). In addition, regarding with availability of the two most common tree species (Table 1), *B. spuria* was frequently associated with *Lysiloma* sp. (67 %). Otherwise, *C. candelaris* (33 %) and the two mentioned *Physcia* species were associated with *S. atomaria* (50 %).

Although specimens of *B. spuria* and *Platythecium grammitis* were not grazed in a tree of *Lysiloma* sp., grazing traces of tree snails were recorded on 100 % of lichen species. The extent of grazing damage to lichen species seems to vary irregularly with the structure of the lichen community on each tree (Fig. 2). Furthermore, among snail feces samples, we identified two lichen species from their spores: *B. spuria* was in

feces of all three tree snail species, and *Coenogonium roumeguerianum* spores were in feces of *P. venusta* and *L. fasciatus*. Four other lichen species were evident from spores but were not identifiable and were recorded as separate entities.

Dispersion of tree snail and corticolous lichen species along the two first NMDS axes (Fig. 3) and the observed frequencies of co-occurrence between snail and lichen species (Fig. 4) documented an association between the snail *H. lucipeta* and the crustaceous lichens *Platythecium grammitis* ( $\chi^2 = 11.6$ ;  $df = 1$ ;  $p < 0.001$ ) and the undetermined lichen species 3 ( $\chi^2 = 100$ ;  $df = 1$ ;  $p < 0.001$ ). Nevertheless, *P. venusta* displayed higher association with trees that carried *B. spuria* ( $\chi^2 = 17.6$ ;  $df = 1$ ;  $p < 0.001$ ) and *C. candelaris* ( $\chi^2 = 36$ ;  $df = 1$ ;  $p < 0.001$ ). The snail *L. fasciatus* significantly associated with the foliose lichens *P. solediosa* ( $\chi^2 = 36$ ;  $df = 1$ ;  $p < 0.001$ ) and *P. aipolia* ( $\chi^2 = 100$ ;  $df = 1$ ;  $p < 0.001$ ). In addition, there was a weakly significant association between *L. fasciatus* and *C. candelaris* ( $\chi^2 = 4$ ;  $df = 1$ ;  $p = 0.045$ ).

#### DISCUSSION

Our finding that lichen species richness per host tree ranged from 1 to 9, with most of the trees bearing between 2 to 6 lichen species, could be potentially explained by substrate associations of lichens which vary according to microhabitat features and forest type. Rosabal *et al.* (2012) did not find any trend in species richness related to changes in tree diameter or bark roughness in a semideciduous forest. By contrast, Rosabal *et al.* (2013) later found that lichen species diversity varied in relation to the diameter of the trunk, pH and phenol concentration of the bark in a montane rain forest. In addition, a constrained distribution of lichen species to certain forest types is expected because lichens often grow under specific micro-niche conditions (Soto-Medina *et al.*, 2012) and their distribution is affected by lichenivorous gastropods (Asplund *et al.*, 2018).

We found that 76.7 % of recorded lichen species were hosted by one tree species, *S. atomaria*, which usually has smooth bark. This morphological trait might increase the possibilities of lichen growth on the bark surface. Similarly, *P. venusta* tree snails were associated with smooth bark at Sardinero locality, Santiago de Cuba province, eastern Cuba (Reyes-Tur and González-Rodríguez, 2003). At this other locality, the lichens *Buellia*, *Lecanora*,

TABLE 1. Data of corticolous lichen species found on 15 host trees of tree snails species (i.e., *Polymita venusta*, *Liguus fasciatus*, *Hemitrochus lucipeta*) and frequency on the two common host tree species from La Rinconada, Jiguaní municipality, Granma, Cuba. [Growth form]: C, Crustaceous; F, Foliose; FB, Foliose with broad lobes; Fr, Fruticose.  
 TABLA 1. Datos de las especies de líquenes cortícolas encontradas sobre 15 árboles hospederos de moluscos arborícolas (i.e., *Polymita venusta*, *Liguus fasciatus*, *Hemitrochus lucipeta*) y su frecuencia sobre las dos especies de árboles más comunes como hospederas, en La Rinconada, municipio Jiguaní, Granma, Cuba. [Forma de crecimiento]: C, Crustáceo; F, Foliáceo; FB, Foliáceo con lóbulos anchos; Fr, Fruticuloso.

Family/ Species	Occurrence frequency (%)	Frequency on <i>Sen-na atomaria</i> (%)	Frequency on <i>Lysiloma</i> sp. (%)
Arthoniaceae			
<i>Arthonia cinnabarina</i> (DC.) Wallr. [C]	20	33.3	0
Arthopyreniaceae			
<i>Bogoriella subfallens</i> (Müll. Arg.) Aptroot & Lücking [C]	6.7	16.7	0
Caliciaceae			
<i>Buellia spuria</i> (Schaer.) Anzi [C]	46.7	33.3	66.7
<i>Dirinaria picta</i> (Sw.) Clem. [F]	13.3	33.3	0
<i>Pyxine</i> sp. [F]	6.7	16.7	0
Chrysotrichaceae			
<i>Chrysothrix candelaris</i> (L.) J.R. Laundon [C]	33.3	33.3	0
Coenogoniaceae			
<i>Coenogonium roumeguerianum</i> (Müll. Arg.) Kalb [C]	20	16.7	0
Graphidaceae			
<i>Graphis adpressa</i> Vain. [C]	13.3	33.3	0
<i>Graphis</i> sp. [C]	6.7	0	0
<i>Platythecium grammitis</i> (Fée) Staiger [C]	20	16.7	66.7
Undetermined species 1 [C]	20	16.7	66.7
Parmeliaceae			
<i>Parmotrema cristiferum</i> (Taylor) Hale [FB]	20	50	0
<i>Parmotrema hypotropum</i> (Nyl.) Hale [FB]	6.7	16.7	0
<i>Parmotrema praesorediosum</i> (Nyl.) Hale [FB]	6.7	16.7	0
<i>Platismatia</i> sp. [F]	13.3	33.3	0
Physciaceae			
<i>Physcia aipolia</i> (Ehrh. ex Humb.) Fűrnr. [F]	26.7	50	0
<i>Physcia americana</i> G. Merrill [F]	6.7	16.7	0
<i>Physcia solediosa</i> (Vain.) Lynge [F]	33.3	50	0
Pyrenulaceae			
<i>Pyrenula leucostoma</i> Ach. [C]	6.7	0	0
Ramalinaceae			
<i>Ramalina peruviana</i> Ach. [Fr]	6.7	0	0
Strigulaceae			
<i>Strigula griseonitens</i> R.C. Harris [C]	13.3	16.7	33.3
Trypetheliaceae			
<i>Astrothelium subcatervarium</i> (Malme) Aptroot & Lücking [C]	6.7	16.7	0
<i>Trypethelium eluteriae</i> Spreng. [C]	6.7	16.7	0
Unknown Family			
Undetermined species 2 [C]	6.7	0	0
Undetermined species 3 [C]	6.7	0	33.3
Undetermined species 4 [F]	6.7	0	0
Undetermined species 5 [F]	6.7	0	0
Undetermined species 6 [C]	6.7	16.7	0
Undetermined species 7 [F]	6.7	16.7	0
Undetermined species 8 [C]	6.7	0	0




FIGURE 3. Non-metric multidimensional scaling (NMDS) plot of the corticolous lichen community (green circles) in relation with three tree snails species (red symbols) from La Rinconada, Jiguani municipality, Granma, Cuba. Lichen species names: Art\_cin = *Arthonia cinnabarina*, Ast\_sub = *Astrothelium subcatervarium*, Bog\_sub = *Bogoriella subfaellens*, Bue\_spu = *Buellia spuria*, Chr\_can = *Chrysothrix candelaris*, Coe\_rou = *Coenogonium roumeguerianum*, Dir\_pic = *Dirinaria picta*, Gra\_adp = *Graphis adpressa*, Par\_cri = *Parmotrema cristiferum*, Phy\_ame = *Phycia aipolia*, Phy\_sor = *Phycia solediosa*, Plat\_gra = *Platythecium grammitis*, Pyr\_leu = *Pyrenula leucostoma*, Ram\_per = *Ramalina peruviana*, Stri\_gris = *Strigula griseonitens*, Try\_elu = *Trypethelium eluteriae* var. *citrinum*, sp 1 = Undetermined species 1, sp 2 = Undetermined species 2, sp 3 = Undetermined species 3, sp 4 = Undetermined species 4, sp 5 = Undetermined species 5, sp 6 = Undetermined species 6, sp 7 = Undetermined species 7, sp 8 = Undetermined species 8, Tree snail species names: Hem\_luc = *Hemitrochus lucipeta*, Lig\_fas = *Liguus fasciatus*, Pol\_ven = *Polymita venusta*.

FIGURA 3. Representación gráfica del Escalamiento multidimensional no métrico (EMDN) de la comunidad de líquenes cortícolas (círculos verdes) relacionada con tres especies de moluscos arborícolas (símbolos rojos) en La Rinconada, municipio Jiguani, Granma, Cuba.

*Opegrapha* and *Graphis* were frequent. In general, it seems that Cuban tree snails such as *Polymita* or *Liguus* are associated with smooth-barked trees. There is often higher crustose lichen abundance on smooth bark (Díaz-Piferrer, 1961; Milera and Martínez, 1987). For *P. muscarum*, Bidart (1997) found an abundance of fungi such as *Aspergillus* sp. and *Pericoma* sp. in the most used host trees (*Eugenia* sp. and *Lysiloma* sp.). Therefore, crustose lichens and non-lichenized fungi could play an important role in the diet of *Polymita* species.

At our Rinconada site we found a significant snail-specific association with certain corticolous lichen species. The species *L. fasciatus* showed a highly significant association with two foliose species frequently associated with *S. atomaria* tree species, *P. solediosa* and *P. aipolia*. By contrast, the tree snail *H. lucipeta*, restricted to the

host *Lysiloma* sp., seems to be associated with the crustose lichens *P. grammitis* and another unknown species. Interestingly, *P. venusta* displayed higher associations with the crustose lichens *B. spuria* and *C. candelaris*, when these were living on *Lysiloma* sp. and *S. atomaria*, respectively. These species-specific association patterns suggest niche segregation by diet as reported by Baur *et al.* (1992, 1994) in two European snails, *Chondrina clienta* (Westerlund, 1883) and *Balea aspersa* (Linnaeus, 1758). They reported a difference between grazed lichen species and they proposed, as an alternative explanation, that divergence of niche diminishes the interspecific competition.

We related our new findings to the phenomenon that snail behavior on tree trunks is integrated with defecation activities. We found spores of six lichen species in tree snails feces, with




FIGURE 4. Frequency of co-occurrence of tree snail and lichen species with significant association on host trees from La Rinconada, Jiguaní municipality, Granma, Cuba.

FIGURA 4. Frecuencia de coincidencia de las especies de líquenes y moluscos arborícolas asociadas significativamente, sobre árboles hospederos en La Rinconada, municipio Jiguaní, Granma, Cuba.

the highest frequencies for *C. roumeguerianum* and *B. spuria* spores in all the studied tree snails. *Polymita venusta* is associated with trees with *B. spuria* as apparently this lichen is an important part of its diet. For *P. venusta*, Reyes-Tur (2004) reported that defecation was a frequent behaviour at the beginning of nocturnal activity. During and immediately after defecation these snails rubbed different parts of their soft bodies with their fresh feces. During daytime most of these individuals (70 %) returned to the sites where they rubbed feces the night before. Hence, in agreement with suggestion of Cook (2001) for other land snail species, it could be possible that *P. venusta* labels suitable feeding and resting sites through substances present in the fecal components (e.g., spores). The evidence of spore diversity within feces and grazed traces both revealed that most of recorded lichen species were consumed by two tree snails, *P. venusta* and *L. fasciatus*. This suggests that at least these two tree snails are generalist herbivores despite spatial association for habitats with particular lichen species.

In conclusion, we described associations between three species of Cuban tree snails and epiphytic lichens on their host trees. Most of the tree snail and lichen species were associated with *S. atomaria*, a tree species that usually has smooth bark. Further, there was not only a significant snail species-specific association for habitats with particular lichen species but also our findings suggested that two species of studied

snails were generalist herbivores. However, in order to provide a thorough account of the patterns of snail-lichen association, more replication is needed to increase the sample size. In addition, we need to increase our knowledge and understanding of the relative palatability of the different lichen species to the three species of tree snail.

**ACKNOWLEDGMENTS.** We are grateful to the Capote-Danet family, Ianna Benavides-Oro and Israelio Marrero for logistic support in the field work. We also thank Mario J. Gordillo-Pérez, Elena Fornet-Balbina, Alena Reyes-Fornet and Nilia Cuéllar-Araújo for remarks on an early draft. Camila Bosch-Díaz helped with figures. Yenisey Rivillo Góngora gave us taxonomic information on some plant species. Yngvar Gauslaa, Steffen Boch, Alison C. Dibble, John A. Allen and three anonymous referees provided very useful writing assistance, comments and suggestions to improve the manuscript. The work was partially financed by VLIRUOS project and we also thank Tino Sauer and the NABU agency for their general support.

## REFERENCES

- Aptroot, A., R. Lücking, H. Sipman, L. Umaña, and J. Chávez. 2008. Pyrenocarpous lichens with bitunicate asci: A first assessment of the lichen biodiversity inventory in Costa Rica. *Bibliotheca Lichenologica* 97: 1-162.
- Asplund, J. and Y. Gauslaa. 2008. Mollusc grazing limits growth and early development of the old forest

- lichen *Lobaria pulmonaria* in broad leaved deciduous forests. *Oecologia* 155: 93-99.
- Asplund, J., Y. Gauslaa, and S. Merinero. 2016. The role of fungal parasites in tri-trophic interactions involving lichens and lichen-feeding snails. *New Phytologist* 211: 1352-1357.
- Asplund, J., P. Larsson, S. Vatne, and Y. Gauslaa. 2010. Gastropod grazing shapes the vertical distribution of epiphytic lichens in forest canopies. *Journal of Ecology* 98: 218-225.
- Asplund, J., O. V. Strandin, and Y. Gauslaa. 2018. Gastropod grazing of epiphytic lichen-dominated communities depends on tree species. *Basic and Applied Ecology* 32: 96-102.
- Asplund, J. and D. A. Wardle. 2013. The impact of secondary compounds and functional characteristics on lichen palatability and decomposition. *Journal of Ecology* 101: 689-700.
- Asplund, J. and D. A. Wardle. 2017. How lichens impact on terrestrial community and ecosystem properties. *Biological Reviews* 92: 1720-1738.
- Baur, A., B. Baur, and L. Fröberg. 1992. The effect of lichen diet on growth rate in the rock-dwelling land snails *Chondrina clienta* (Westerlund) and *Balea perversa* (Linnaeus). *Journal of Molluscan Studies* 58: 345-347.
- Baur, A., B. Baur, and L. Fröberg. 1994. Herbivory on calcicolous lichens: different food preferences and growth rates in two co-existing land snails. *Oecologia (Berl.)* 98: 313-319.
- Baur, B., L. Fröberg, and A. Baur. 1995. Species diversity and grazing damage in a calcicolous lichen community on top of stone walls in Öland, Sweden. *Annales Botanici Fennici* 32: 239-250.
- Baur, B., L. Fröberg, A. Baur, R. Guggenheim, and M. Haase. 2000. Ultrastructure of snail grazing damage to calcicolous lichens. *Nordic Journal of Botany* 20: 119-128.
- Bidart, L. 1997. Ecología de *Polymita muscarum* Lea, 1834 (Gastropoda: Xanthonychidae) en la provincia de Holguín, Cuba. [Unpublished]. MSc Thesis. Instituto de Ecología y Sistemática, La Habana, Cuba. 50 pp.
- Boch, S., M. Fischer, and D. Prati. 2015. To eat or not to eat—Relationship of lichen herbivory by snails with secondary compounds and field frequency of lichens. *Journal of Plant Ecology* 8: 642-650.
- Boch, S., D. Prati, and M. Fischer. 2016. Gastropods slow down succession and maintain diversity in cryptogam communities. *Ecology* 97: 2184-2191.
- Brodo, I. M., S. D. Sharnoff, and S. Sharnoff. 2001. *Lichens of North America*. Yale University Press, New Haven and London. 795 pp.
- Cameron, R. 2016. *Slugs and snails*. Harper Collins Publishers, London. 508 pp.
- Capote, R. P. and R. Berazaín. 1984. Clasificación de las formaciones vegetales de Cuba. *Revista Jardín Botánico Nacional* 5(2): 27-75.
- Clarke, K. R. and R. N. Gorley. 2006. PRIMER v6: user manual/tutorial. PRIMER-E, Plymouth.
- Coker, P. D. 1967. Damage to lichens by gastropods. *Lichenologist* 3: 428-429.
- Cook, A. 2001. Behavioural ecology: On doing right thing, in the right place at the right time. Pp. 447-487. In: *The biology of terrestrial molluscs* (G. M. Barker, Ed.). CABI Publishing, Wallingford, Nueva York, EUA. 558 pp.
- Crous, P. W., W. Gams, J. A. Stalpers, V. Robert, and G. Stegehuis. 2004. MycoBank: an online initiative to launch mycology into the 21<sup>st</sup> century. *Studies in Mycology* 50: 19-22.
- Curry, J. P. 1994. *Grassland invertebrates. Ecology, influence on soil fertility and effects on plant growth*. Chapman & Hall, London. 437 pp.
- Díaz-Piferrer, M. 1961. Feeding habitat of a Cuban tree snail (*Polymita muscarum* Lea). *Caribbean Journal of Science* 1: 123-132.
- Fröberg, L., A. Baur, and B. Baur. 1993. Differential herbivore damage to calcicolous lichens by snails. *Lichenologist* 25: 83-85.
- Fröberg, L., A. Baur, and B. Baur. 2006. Field study on the regenerative capacity of three calcicolous lichen species damaged by snail grazing. *Lichenologist* 38: 491-493.
- Fröberg, L., P. Stoll, A. Baur, and B. Baur. 2011. Snail herbivory decreases cyanobacterial abundance and lichen diversity along cracks of limestone pavements. *Ecosphere* 2.
- González-Guillén, A. 2014. *Polymita the most beautiful land snail of the world*. Fundcraft Publishing, Collierville, Tennessee. 359 pp.
- González-Guillén, A., F. Krull, and L. Álvarez-Lajonchere Ponce de León. 2018. *Liguus the flamboyant tree snails*. Fundcraft Publishing, Collierville, Tennessee. 244 pp.
- Hatzioannou, M., N. Eleutheriadis, and M. Lazaridou-Dimitriadou. 1994. Food preferences and dietary overlap by terrestrial snails in Logos area (Edessa, Macedonia, Northern Greece). *Journal of Molluscan Studies* 60: 331-341.
- Hawksworth, D. L. and R. Lücking. 2017. Fungal diversity revisited: 2.2 to 3.8 million species. *Microbiology Spectrum* 5(4): FUNK-0052-2016.
- Hernández-Quinta, M., L. Álvarez Lajonchere Ponce de León, D. Martínez Borrego, D. Maceira Filgueira, A. Fernández Velázquez, and J. Espinosa Sáez. 2017. Moluscos terrestres y dulceacuícolas. Pp. 168-195. In: *Diversidad biológica de Cuba: métodos de inventario, monitoreo y colecciones biológicas* (C. A. Mancina and D. D. Cruz, Eds.). Editorial AMA. La Habana, 502 pp.
- Jongman, R. H. G., C. J. F. ter Braak, and O. F. R. van Tongeren. 2001. *Data analysis in community and landscape ecology*. Cambridge University Press, Cambridge. 299 pp.

- Lawrey, J. D. 1983. Lichen herbivore preference: a test of two hypotheses. *American Journal of Botany* 70: 1188-1194.
- Lücking, R. and A. Bernecker-Lücking. 2000. Lichen feeders and lichenicolous fungi: do they affect dispersal and diversity in tropical foliicolous lichen communities? *Ecotropica* 6: 23-41.
- Lücking R., M. Dal-Forno, M. Sikaroodi, P. M. Gillevet, F. Bungartz, B. Moncada, A. Yáñez-Ayabaca, J. L. Chaves, L. F. Coca, and J. D. Lawrey. 2014. A single macrolichen constitutes hundreds of unrecognized species. *Proceedings of the National Academy of Sciences of the United States of America* 111: 11091-11096.
- Milera, J. F. and J. R. Martínez. 1987. *Polymita*. Editorial Científico Técnica, La Habana. 70 pp.
- Minter, D. W., M. Mena-Portales, J. Rodríguez-Hernández, H. Iglesias-Brito, M. Camino-Vilaró, and A. Mercado-Sierra. 2002. Mapas de Distribución Computarizados de los Hongos del Caribe. Electronic publication on CD, containing 11950 HTML-format pages and 11756 GIF-format maps. PDMS Publishing, Isleworth.
- Minter, D. W., M. Rodríguez-Hernández, and J. Mena-Portales. 2001. Fungi of the Caribbean. An annotated checklist. Middlesex, UK. 946 pp.
- Moberg, R. 1990. The lichen genus *Physcia* in Central and South America. *Nordic Journal Botany* 10: 319-342.
- Peake, J. F. and P. W. James. 1967. Field and study notes. *Lichenologist* 3: 425-428.
- Pilsbry, H. P. 1946. Land Mollusca of North American (north of Mexico). *Monographs of the Academy of Natural Sciences of Philadelphia* 3: 1-520.
- Pluntke, M. 1984. Die Flechtenflora Kubas (Flora Lichenum Cubensis). *Terrestrische Ökologie. Sonderheft 4. Universitäts und Landesbibliothek. Sachsen-Anhalt.* 157 pp.
- Price, M. R., R. O'Rorke, A. S. Amend, and M. G. Hadfield. 2016. Diet selection at three spatial scales: Implications for conservation of an endangered Hawaiian tree snail. *Biotropica* 49: 130-136.
- Quinn, G. P. and M. J. Keough. 2002. *Experimental design and data analysis for biologists*. Cambridge University Press, Cambridge. 537 pp.
- Reyes-Tur, B. 2004. Ecología y biología reproductiva de *Polymita venusta* (Lea, 1834). [Unpublished]. Tesis de Doctorado. Universidad de La Habana. Cuba. 100 pp.
- Reyes-Tur, B. and A. González-Rodríguez. 2003. Relación planta-animal del molusco terrestre cubano *Polymita venusta* (Stylommatophora: Helminthoglyptidae) en Sardinero, Santiago de Cuba, Cuba. *Revista Biología* 17: 134-142.
- Robert, V., G. Stegehuis, and J. Stalpers. 2005. The MycoBank engine and related databases. <http://www.mycobank.org/> (accessed 2 November 2018).
- Robert, V., D. Vu, A. Ben Hadj Amor, N. van de Wiele, C. Brouwer, B. Jabas, S. Szoke, A. Dridi, M. Triki, S. ben Daoud, O. Chouchen, L. Vaas, A. de Cock, J. A. Stalpers, D. Stalpers, G. J. M. Verkley, M. Groenewald, F. Borges dos Santos, G. Stegehuis, W. Li, L. Wu, R. Zhang, J. Ma, M. Zhou, S. Pérez Gorjón, L. Eurwilaichitr, S. Ingsriswang, K. Hansen, C. Schoch, B. Robbertse, L. Irinyi, W. Meyer, G. Cardinali, D. L. Hawksworth, J. W. Taylor, and P. W. Crous. 2013. [MycoBank gearing up for new horizons](#). *IMA Fungus* 4: 371-379.
- Rollo, C. D. 1988. The feeding of terrestrial slugs in relation to food characteristics, starvation, maturation and life history. *Malacologia* 28: 29-39.
- Rosabal, D., A. R. Burgaz, A. Altamirano, and G. Aragón. 2012. Differences in diversity and reproductive strategies of corticolous lichens between interior and edge of the Monte Barranca semideciduous forest, Santiago de Cuba. *Bryologist* 115: 333-340.
- Rosabal, D., A. R. Burgaz, and O. J. Reyes. 2013. Substrate preferences and phorophyte specificity of corticolous lichens on five tree species of the montane rainforest of Gran Piedra, Santiago de Piedra, Santiago de Cuba. *Bryologist* 116: 113-121.
- Rosabal, D., A. R. Burgaz, and O. J. Reyes. 2016. Analizando la diversidad beta en ensamblajes de líquenes en un gradiente vertical sobre cinco especies de forófitos en la pluvisilva montana de la Gran Piedra, Cuba. *Botanica Complutense* 40: 31-40.
- Shachak, M., C. G. Jones, and Y. Granot. 1987. Herbivory in rock and the weathering of a desert. *Science* 236: 1098-1099.
- Soto-Medina, E., R. Lücking, and A. Bolaños Rojas. 2012. Especificidad de forófito y preferencias microambientales de los líquenes cortícolas en cinco forófitos del bosque premontano de finca Zingara, Cali, Colombia. *Revista Biología Tropical* 60: 843-856.
- Speiser, B. 2001. Food and feeding. Pp. 259-288. In: *The biology of terrestrial molluscs* (G. M. Barker, Ed.). CABI Publishing. Wallingford, Nueva York, EUA. 558 pp.
- StatSoft, Inc. 2003. Statistica versión 6.1. Available in <http://www.statsoft.com/> (accessed 20 December 2006).
- Torre, C. de la. 1950. El género *Polymita*. *Memorias de la Sociedad Cubana de Historia Natural "Felipe Poey"* 20: 1-20.
- Vatne, S., T. Solhøy, J. Asplund, and Y. Gauslaa. 2010. Grazing damage in the old forest lichen *Lobaria pulmonaria* increases with gastropod abundance in deciduous forests. *Lichenologist* 42: 615-619.
- Wirth, M. and M. E. Hale. 1978. Morden-Smithsonian expedition to Dominica: The lichens (Graphidaceae). *Smithsonian contributions to Botany* 40: 1-64.

Recibido: 9 de enero de 2020/aceptado: 15 de mayo 2020