




Surveying lichen diversity in forests: A comparison of expert mapping and eDNA metabarcoding of bark surfaces

Lukas Dreyling^{1,2}, Steffen Boch³, H. Thorsten Lumbsch⁴, Imke Schmitt^{1,2}

1 Senckenberg Biodiversity and Climate Research Centre (SBIK-F), Frankfurt am Main, Germany

2 Goethe University Frankfurt, Institute of Ecology, Evolution and Diversity, Frankfurt am Main, Germany

3 WSL Swiss Federal Institute for Forest, Snow and Landscape Research, Birmensdorf, Switzerland

4 Collections, Conservation, and Research, The Field Museum, Chicago, IL 60605-2496, USA

Corresponding authors: Lukas Dreyling (Lukas.Dreyling@gmx.net); Imke Schmitt (imke.schmitt@senckenberg.de)

Abstract

Lichens are an important part of forest ecosystems, contributing to forest biodiversity, the formation of micro-niches and nutrient cycling. Assessing the diversity of lichenised fungi in complex ecosystems, such as forests, requires time and substantial skills in collecting and identifying lichens. The completeness of inventories thus largely depends on the expertise of the collector, time available for the survey and size of the studied area. Molecular methods of surveying biodiversity hold the promise to overcome these challenges. DNA barcoding of individual lichen specimens and bulk collections is already being applied; however, eDNA methods have not yet been evaluated as a tool for lichen surveys. Here, we assess which species of lichenised fungi can be detected in eDNA swabbed from bark surfaces of living trees in central European forests. We compare our findings to an expert floristic survey carried out in the same plots about a decade earlier. In total, we studied 150 plots located in three study regions across Germany. In each plot, we took one composite sample based on six trees, belonging to the species *Fagus sylvatica*, *Picea abies* and *Pinus sylvestris*. The eDNA method yielded 123 species, the floristic survey 87. The total number of species found with both methods was 167, of which 48% were detected only in eDNA, 26% only in the floristic survey and 26% in both methods. The eDNA contained a higher diversity of inconspicuous species. Many prevalent taxa reported in the floristic survey could not be found in the eDNA due to gaps in molecular reference databases. We conclude that, currently, eDNA has merit as a complementary tool to monitor lichen biodiversity at large scales, but cannot be used on its own. We advocate for the further development of specialised and more complete databases.

Key words: Assessment, biodiversity, bioindicators, conservation, databases, floristic survey, identification, inventory, metabarcoding, monitoring



Academic editor: Pradeep Divakar

Received: 18 December 2023

Accepted: 10 May 2024

Published: 21 June 2024

Citation: Dreyling L, Boch S, Lumbsch HT, Schmitt I (2024) Surveying lichen diversity in forests: A comparison of expert mapping and eDNA metabarcoding of bark surfaces.

MycoKeys 106: 153–172. <https://doi.org/10.3897/mycokeys.106.117540>

Copyright: © Lukas Dreyling et al.

This is an open access article distributed under terms of the Creative Commons Attribution

License (Attribution 4.0 International – CC BY 4.0).

Introduction

Lichens are important components of biodiversity in forest ecosystems, where they form epiphytic communities in the canopy (Ellis 2012) and on tree trunks (Hofmeister et al. 2016). In central European forests, lichens and their symbionts are characteristic taxa of the bark surface community (Baldrian 2017; Dreyling

et al. 2022; Hofmann et al. 2023). In fact, the bark of trees has been proposed as an important part of the forest microbiome and sustains a high microbial biomass especially if lichens are present (Baldrian 2017). Temperate forests harbour more than a hundred, often several hundred species of lichen-forming fungi (Jüriado et al. 2003; Coppins and Coppins 2005; Boch et al. 2013; Löhmus and Löhmus 2019). The lichen communities contribute to forest ecosystem function by retaining water (Van Stan and Pypker 2015), cycling minerals and nutrients (Pike 1978; Reiners and Olson 1984; Knops et al. 1991, 1996; Campbell et al. 2010), being part of the food web and providing habitat and micro-niches for other organisms (reviewed in Ellis (2012) and Asplund and Wardle (2017)). Forest lichen communities respond to abiotic environmental changes (Miller et al. 2018; Łubek et al. 2021), as well as to forest management (Nascimbene et al. 2013; Boch et al. 2021). Some species can be used as indicators to monitor the effects of anthropogenic pollutants (Fрати and Brunialti 2023). These are important reasons to survey and monitor lichen biodiversity in forests.

The assessment of lichen biodiversity can be challenging, even for taxonomic experts (Vondrák et al. 2016). Since species identification of lichenised fungi often relies only on few morphological characters (Crespo and Lumbsch 2010), considerable expertise is necessary and often requires specimen collection for ex situ identification, for example, through microscopy or chemical tests (Wright et al. 2019). As a result, the outcomes of lichen surveys are highly dependent on the training of collectors (Giordani et al. 2009). Additionally, lichen-forming fungi are a group with high potential for cryptic diversity that cannot be distinguished in the field (Crespo and Lumbsch 2010; Altermann et al. 2014).

Molecular markers are useful complementary tools to aid the identification of lichenised and non-lichenised fungi (Lücking et al. 2020). Especially, the ITS barcode marker is increasingly used for species identification and species delimitation of lichenised fungi (Schoch et al. 2012; Bradshaw et al. 2020). It has been applied to assess species diversity within geographic regions (Kelly et al. 2011), as well as within taxonomic groups, for example, Parmeliaceae (Divakar et al. 2016). While ITS barcoding works well for the majority of species, it has limitations in some taxonomic groups, for example, the genus *Cladonia*, which seems to be lacking a sufficient barcode gap at least in some species complexes (Pino-Bodas et al. 2013; Marthinsen et al. 2019) or members of Graphidaceae and Pertusariaceae, which do not amplify reliably with common ITS primers, so that their ITS is not used in multi-locus phylogenies (Schmitt and Lumbsch 2004; Rivas Plata et al. 2013) and they remain under-represented in DNA databases. A small number of studies have attempted to identify species and characterise the lichen community by metabarcoding bulk specimen collections (Wright et al. 2019; Henrie et al. 2022). They found that this method produces comparable results between minimally trained and expert collectors and thus potentially reduces the need for extensive training (Wright et al. 2019; Henrie et al. 2022). Furthermore, they concluded that metabarcoding surveys could enable a more efficient sampling over a larger spatial extent (Wright et al. 2019).

Biodiversity assessments using environmental DNA (eDNA) allow species-level identification from DNA present in environmental samples, such as water, soil or air (Taberlet et al. 2012; Yoccoz 2012). In comparison to bulk metabarcoding, this method has additional advantages, such as targeting a broader range of taxa (Taberlet et al. 2012) and being non-invasive, i.e. not

requiring destruction of specimens (Deiner et al. 2017). Despite some drawbacks, in particular due to incomplete databases and primer bias (Bellemain et al. 2010; Keck et al. 2022), eDNA has shown great potential for biodiversity assessments (Shirouzu et al. 2016; Frøslev et al. 2019). When eDNA and conventional methods were compared, species overlap was variable depending on the taxonomic group, but eDNA always identified taxa that were not picked up with other methods (Cordier et al. 2021). In a meta-analysis, eDNA was found to detect more species in general and significantly more rare species, exhibiting higher accuracy and efficiency, while being less costly than conventional biodiversity assessments (Fediajevaite et al. 2021).

In this study, we analyse the utility of eDNA – obtained from bark surfaces of tree trunks at breast height – to assess the diversity of lichen communities in central European forests. In previous studies, we have generated datasets of entire fungal communities associated with bark surfaces, based on ITS metabarcoding (Dreyling et al. 2022, 2024; Hofmann et al. 2023). Here, we use only the fraction of lichenised fungi from these datasets. The sampling sites are 150 specific plots located within the Biodiversity Exploratories (Fischer et al. 2010) in northern, central and southern Germany. We compare the results to a previous floristic survey carried out in the same plots (Boch et al. 2021). Specifically, we address the following questions: I. Which species of lichenised fungi can be identified from environmental samples via eDNA metabarcoding? II. What are the differences to the diversity obtained through an expert survey? III. Is eDNA metabarcoding a reliable tool to survey lichen diversity in forests?

Material and methods

Study sites

We surveyed communities of forest-dwelling lichen species in 150 plots, located in three regions, within the Biodiversity Exploratories framework (Fischer et al. 2010). The regions mark a south-west to north-east gradient across Germany and differ in their climate and topography. The plots within the regions were further selected along a gradient of the anthropogenic impact, i.e. forest management intensity (Fischer et al. 2010; Boch et al. 2021) and are representative for central European forest ecosystems. The South-West region is on average 2 °C colder than the North-East (6.5 °C vs. 8.5 °C) and experiences approximately twice the amount of precipitation (700–1000 mm vs. 500–600 mm; Fischer et al. (2010)). In addition, they differ in their tree species composition, with beech trees (*Fagus sylvatica*) as the dominant species in most plots across the regions. Some plots are dominated by coniferous trees, specifically Norway spruce (*Picea abies*) in the South-West region and Scots pine (*Pinus sylvestris*) in the North-East region (Schall and Ammer 2018). For both, eDNA sampling and classical lichen mapping, we surveyed a 20 m × 20 m subplot within the established 100 m × 100 m experimental plots.

eDNA sampling and processing

In each plot, we collected eDNA samples from the bark surface of six trees of the respective dominant species in May 2021. The six individual tree samples were pooled into one composite sample per plot. Since we had previously

shown large community differences between tree sizes (Dreyling et al. 2022), we included two trees each of large (> 30 cm diameter at 150 cm height), medium (15–30 cm) and small (5–15 cm) size in each sample. If this type of sampling was not possible, we included additional trees of the size class that best represented the forest in the immediate surrounding. To sample the bark surface eDNA, we moistened the bark surface with sterile water and then used a nylon-flocked medical swab (FLOQSwabs™, Copan, Brescia, Italy) to collect the bark surface biofilm (Fig. 1). We swabbed around the full tree trunk at approximately 150 cm height from the forest floor, excluding large patches of bryophytes to avoid bias due to the amplification of plant ITS, but explicitly including all other epiphytic organisms. The material collected with the swabs was fixed with 5 ml nucleic acid preservation (NAP) buffer (Camacho-Sanchez et al. 2013) in 15 ml tubes directly after sampling and placed on ice in the field. Afterwards, the samples were stored at 4 °C until DNA extraction in the following week.



Figure 1. Sampling procedure: We moistened the tree trunk on all sides at breast height and swabbed the bark surface in a zigzagging motion along a 10 cm wide band around the tree trunk. The swabbed area included smooth bark surfaces and crevices, as well as epiphytic organisms, if they were present.

A detailed description of the DNA extraction and bioinformatic processing of sequencing reads is given in Dreyling et al. (2022). In brief: We extracted DNA, from samples as well as three extraction blanks, using an extraction kit (Quick-DNA Fecal/Soil Microbe Microprep, Zymo Research Europe GmbH, Freiburg, Germany) with an additional step ensuring liberation of material from the swab. Targeting the ITS2 region, we subsequently amplified the fungal DNA in triplicate, using the universal primer pair fITS7 (GTGARTCATCGAATCTTTG) (Ihrmark et al. 2012) and ITS4 (TCCTCCGCTTATTGATATGC) (White et al. 1990). PCR reactions also included negative controls (without sample material) and multiplex controls (empty wells). We cleaned the amplicons via a magnetic bead protocol (MagSI-NGSPREP Plus, magtivio B.V., Geelen, Netherlands) and measured DNA concentration through fluorometry (Qubit dsDNA HS assay on a Qubit 3.0, Thermo Fisher Scientific, MA, United States), before equimolar pooling. The library preparation and Illumina sequencing (MiSeq 2 × 300 bp paired-end) was carried out by Fasteris SA (Plan-les-Ouates, Switzerland) according to their MetaFast Protocol, designed to avoid additional PCR bias.

We used Cutadapt (v3.3; Martin (2011)) to demultiplex the obtained sequencing reads and DADA2 (Callahan et al. 2016) to infer Amplicon Sequencing Variants (ASVs). Taxonomy was assigned against the Martin7 database (Vondrák et al. 2023) using a local BLAST (Altschul et al. 1990) search. Assignments were kept if the “percent identity” was higher than 97%. Additionally, we used the UNITE database (Abarenkov et al. 2022, Version 9.0, incl. non-fungal eukaryotic DNA as outgroups) and the NCBI nucleotide database (Sayers et al. 2022, percent identity > 97%) to assign additional taxonomy to the ASVs that could not be assigned with the Martin7 database. We used FUNGuild (Nguyen et al. 2016) to assign information on the functional guild to the additionally assigned fungal ASVs and filtered the dataset to only contain ASVs which were classified as lichenised fungi. All scripts on the bioinformatic processing, as well as the analysis, are available at Github at https://github.com/LukDrey/eDNA_lichen_survey.

Traditional floristic survey

The floristic survey was carried out in 2007 and 2008 and recorded occurrences of lichenised fungi in over 600 plots of the Biodiversity Exploratories (Boch et al. 2021), including the 150 plots of the eDNA sampling. The survey covered a comparable area of 20 m × 20 m around the plot centre, which was not fully identical, but always spatially close to the sampling area of the eDNA survey. All lichens occurring on bark (up to 2.5 m height), rocks, deadwood and soil were recorded. We did not limit the survey time per plot due to strongly varying environmental heterogeneity amongst plots. Most specimens were identified in situ, except when microscopic or chemical characters had to be assessed.

Comparison of the two methods

A number of taxonomic changes took place in the approximately 13 years between the two surveys. We accommodated for these developments by harmonising the two species lists and adopting the names accepted as the current names in MycoBank (Crous et al. 2004; Robert et al. 2013). Additionally, we

only included species from the floristic dataset, which were recorded as epiphytes in the 150 experimental plots, thus excluding species collected from rocks, deadwood and soil. A list of all species is provided as Suppl. material 1.

To allow for comparisons between the two methods, we transformed the read counts obtained through the eDNA metabarcoding to presence-absence data. Using the two presence-absence datasets, we compared the two methods and assessed the diversity and species richness found with each method. Furthermore, we calculated the number of plots in which a species was found. Finally, we selected five species to visualise geographical occurrence patterns, based on the two different assessment methods.

Results and discussion

In total, we found 167 species of lichenised fungi in the two surveys. The eDNA method found 123 species, while the traditional floristic survey recorded 87 species (Fig. 2). With the eDNA method, we found 80 species that were not found via the traditional survey methods, while with the floristic survey, we found 44 species not detected by the eDNA method. The higher number of taxa identified from the eDNA is congruent with bulk-specimen sequencing studies from other ecosystems (Wright et al. 2019). Interestingly, only 26% of the total taxa were shared between both methods, likely due to the number of small, inconspicuous genera, such as *Micarea* (Launis et al. 2019), that were only found with the eDNA (Fig. 3). The overlap between the two methods is similar to what has previously been reported for comparisons of eDNA to fruiting body surveys of forest fungi (Shirouzu et al. 2016; Frøslev et al. 2019).

In our study, several species are detected exclusively or predominantly by either of the two methods. Additionally, even the most common species are not necessarily detected by both methods. For example, out of the five most common species (Fig. 3), two were not found in the traditional floristic study,




Figure 2. Number and proportion of species of lichenised fungi found either in eDNA metabarcoding or floristic survey or in both methods.

Scoliciosporum sarothamni and *Micarea czarnotae*. This result is especially striking for *S. sarothamni* which was found in 146 of 150 plots via eDNA metabarcoding, but was not identified in the floristic study. This species is very small and thus hard to find and distinguish (Kowalewska and Kukwa 2003; Dymytrova 2011). Therefore, it is plausible that it has been overlooked or simply grouped with other taxa, such as its sister species *Scoliciosporum chlorococcum*, in the floristic survey. However, *S. chlorococcum* was also highly prevalent in the eDNA study (121 occurrences), but rarely found in the floristic dataset (11 occurrences). A potential reason for *Scoliciosporum* being less prevalent in the floristic dataset is that both species have a greenish thallus and are often occurring within dominant green algal colonies, making them hard to recognise with traditional methods, especially when sterile.

Other taxa, commonly found in the eDNA metabarcoding dataset, had not been formally described at the time of the floristic survey. For example, both *Opeltiella rubrisoli* and *Micarea czarnotae* were only described in 2019 (Launis et al. 2019; Liu et al. 2019). *Micarea czarnotae* had previously been included in *M. prasina*, which was also only found in eight plots in the floristic study. In general, the eDNA dataset contained a high number of inconspicuous taxa from genera that are difficult to distinguish, such as *Micarea* (Launis et al. 2019), *Scoliciosporum* (Dymytrova 2011) and *Bacidina* (Czarnota and Guzow-Krzemińska 2018). Consistent with our findings, other studies have previously reported that eDNA was superior in revealing hidden diversity for fungi (Shirouzu et al. 2016), including lichen-forming fungi (Wright et al. 2019). An additional advantage of the eDNA approach might be the detection of taxa not directly occurring on the sampled substrate itself, for example, from propagules (Wright et al. 2019; Henrie et al. 2022).

The floristic dataset also includes numerous taxa which were not identified in the eDNA approach. For example, *Pseudosagedia aenea*, a common species found in the floristic survey (occurring in 104 plots), was not found by the eDNA metabarcoding (Fig. 3), albeit ITS sequences of this taxon are included in the sequence repositories used in this study. One potential reason is that their habitat is outside of the sampled area, for example, in the tree crowns or at the base of the tree. In fact, several species prevalent in the floristic dataset, but not the eDNA, occur in these habitats, for example, *Cladonia coniocraea* at the base of trees (Wirth et al. 2013), *Pseudevernia furfuracea* on branches in the canopy (Kranter et al. 2003) and also *P. aenea* at the stem base (Larsen et al. 2020). Therefore, restricting the eDNA sampling, or any survey, to a single forest substrate is likely insufficient to describe the full lichen diversity (see also Boch et al. (2013)).

Overall, only very few species were found in a similar number of plots with both methods (Fig. 3). The most prevalent species found with both methods were *Coenogonium pineti* and *Lepraria incana*. *Coenogonium pineti* was found in 119 plots in the floristic study and in 80 plots in the eDNA dataset. The prevalence of *L. incana* was even more similar, being found in 106 plots with floristic and 113 plots with eDNA methods. Although, both species preferentially grow at the base of trees (Lackovičová and Guttová 2005; Larsen et al. 2020), they were also often recorded with the eDNA method. It is possible that the dispersal units of these taxa (ascospores in *C. pineti* and soredia in *L. incana*) are dispersed further up the stem, for example, by wind. Furthermore, snails and slugs may play a role in distributing lichen propagules along the stem (Asplund et al. 2010; Boch et al. 2011).




Figure 3. Most common species of lichenised fungi detected by either method (eDNA metabarcoding or floristic survey). We show taxa, which occurred in at least 25 plots (out of 150) across the three regions.

Another apparent reason for the differences in eDNA and floristic surveys are related to the databases necessary for taxonomic assignment of the metabarcoding reads. Despite large efforts in recent years towards the development of reference databases for fungal taxonomy, like the UNITE database (Abarenkov et al. 2023) or the GlobalFungi project (Větrovský et al. 2020), many gaps remain. In our study, several species, commonly found in the floristic study, have no reference sequences in the UNITE database, including *Diarthonis spadicea* and all species of the genus *Arthonia*. Previous studies have proposed to close the gaps in the reference databases by large scale sequencing of lichen herbarium specimens (Gueidan and Li 2022). Regional databases, for example, for Great Britain and Ireland (Kelly et al. 2011) or part of the western USA (Kerr and Leavitt 2023) were helpful in identifying lichen specimens, based on barcodes or bulk metabarcoding. The recently-published Martin7 database, focusing on central European lichens (Vondrák et al. 2023), greatly improved the results of the present study. It enabled the assignment of taxonomy to over 30 additional ASVs, resulting in 27 additional species compared to an initial assignment using the UNITE database.

Technical issues related to sequencing might be the reason that some species present in the floristic study could not be found in the eDNA assessment, although the ITS sequences are included in the UNITE and Martin7 databases. A search with Primer-BLAST (Ye et al. 2012) revealed that, in some cases, the primer combination used in this study could likely not amplify these species. However, some species, commonly occurring in the floristic study such as *Hypogymnia physodes* or *Pseudevernia furfuracea*, should have been amplified with the current primers, indicating other issues. A potential reason might be PCR biases influencing which taxa or groups are preferentially amplified (Bellemain et al. 2010). For example, shorter DNA fragments are usually amplified more often (Deagle et al. 2006). It is possible that we missed ITS sequences that are longer because they contain introns, a frequent and stochastic feature of the rDNAs of lichen-forming fungi (Simon et al. 2005). Furthermore, the output of the sequencing machine is limited, so that taxa with few copies might not be sequenced (Gloor et al. 2017).

There is a temporal gap of approximately 13 years between the floristic survey and the eDNA sampling, which may explain some of the observed differences, especially with regard to pollution with sulphur dioxide and nitrogen. Sulphur dioxide (SO₂) pollution has been decreasing in western Europe since the 1970s, enabling the return of many species to formerly uninhabitable ecosystems (Rose and Hawksworth 1981; Nash and Gries 2002). Conversely, species that are tolerant to acidic and sulphur-enriched conditions, for example, *Lecanora conizaeoides*, have been reported to decline in central Europe (Nash and Gries 2002; Farkas et al. 2022). In our study, the number of plots, in which *L. conizaeoides* was identified with eDNA in 2021, has halved in relation to the floristic study in 2007/2008. Today, nitrogen pollution is more important in shaping lichen communities than SO₂ pollution (Purvis et al. 2003; Hultengren et al. 2004; Pinho et al. 2008; Gadsdon et al. 2010). Temperate forests experience increased deposition of nitrogen, for example, through ammonia fertilisers or nitric oxides from fuel combustion, and nitrophytic species have increased in the recent past (Carter et al. 2017). In the present study, two species regarded as nitrophytes, *Physcia adscendens* and *P. tenella* (Gadsdon et al. 2010), have been found more frequently in the eDNA sampling than in the earlier floristic survey (Fig. 3). Interestingly, other nitrophytic species, such as *Xanthoria parietina*, *Phaeophyscia orbicularis* or *Candelariella reflexa* (Gadsdon et al. 2010), were found less frequently or not at all in the eDNA sampling (Suppl. material 1). In addition, differences between the floristic and the eDNA survey in lichen diversity and community composition might be because of successional developments or the disruption of such developments by forestry management, leading to changes in forest structure and composition, i.e. changed environmental conditions. Such changes might have been even accelerated by climate change that has been proposed to change lichen diversity and community composition (van Herk et al. 2002; Aptroot 2009; Allen and Lendemer 2016; Nascimbene et al. 2016; Nelsen and Lumbsch 2020).

The three study regions differed considerably in their lichen diversity. In the eDNA metabarcoding survey, the proportion of fungal reads assigned to lichens was highest in the south-western region with approximately 39% of the total fungal reads, 27% in the north-eastern and lowest in the central region with only 14%. On average, lichens accounted for 27% of the total fungal reads. We

observe a similar pattern in the floristic survey, where the highest number of species was also recorded in the south-western region (82 species), followed by the central and the north-eastern region (32 species). Previous studies in the Biodiversity Exploratories found similar relationships between the regions for plants (Klaus et al. 2013) and arthropods (Simons et al. 2014) that are potentially explained by differences in climate, land-use intensity or nutrient availability. In our study, the higher species richness in the South-West region is likely related to the higher annual precipitation (Fischer et al. 2010), which has been shown to positively influence lichen richness (Marini et al. 2011), but also because of the generally lower former SO₂ deposition compared to the other two regions (Umweltbundesamt 2005).

The differences amongst the three study regions are also apparent in the distribution maps of the five example species, *Buellia griseovirens*, *Graphis scripta*, *Lepraria incana*, *Phlyctis argena* and *Physcia adscendens*. These species were chosen as examples because they were amongst the most prevalent species (Fig. 3) found with both methods, but varied in how often they were recorded. In general, the highest number of plots, in which the example species were recorded, were located in the South-West region (Fig. 4A), while they did not occur in most plots of the North-East region (Fig. 4 C). If a lichen species was frequently recorded by both methods, such as *B. griseovirens*, *L. incana* and *P. argena* in the South-West (Fig. 4A), then it was also found by one of the methods in spatially close plots. In general, the lichen records do not follow a clearly distinguishable pattern of spatial clustering within the regions.




Figure 4. Distribution of five example species within the analysed forest plots of the three regions. Shown are occurrence data based on the floristic survey and eDNA metabarcoding. Each map represents one region (Biodiversity Exploratory) **A** South-West (Swabian Alb) **B** Central (Hainich-Dün) **C** North-East (Schorfheide-Chorin). Each circle depicts a 100 m × 100 m forest plot.

The detection of these five lichen species was different between the methods in the each of the regions. Of the five example species, only *L. incana* was consistently found with both methods across the three regions and consequently is one of the most prevalent species we found. *B. griseovirens* and *P. argena* were found more often in the eDNA samples and almost exclusively with eDNA in the Central and North-East regions (Fig. 4B, C). It is tempting to speculate about a northward shift of the distribution of these species considering the time difference between the two studies, which could explain the absence in the Central and North-East regions during the floristic survey. In addition, the considerably decreased pollution in these two regions might have led to the recovery of lichen communities with many species re-colonising such formerly heavily polluted areas (e.g. Gilbert (1992)).

Nevertheless, *Graphis scripta* was rarely found in the eDNA, but recorded across all three regions in the traditional survey. This pattern is likely related to the use of ITS2 as a molecular marker in the eDNA, which has previously shown low amplification rates for the genus *Graphis* (e.g. Kraichak et al. (2019)). Interestingly, *P. adscendens* was found with both methods in the South-West, purely with the eDNA in the Central and only via floristic survey in the North-East region.

Conclusions

In its current form, eDNA metabarcoding cannot be used as a stand-alone tool to survey epiphytic lichen diversity. However, it can serve as a valuable complementary tool, similarly to studies from many other taxonomic groups (Beng and Corlett 2020; Fediajevaite et al. 2021). In the long run, with more correct and more complete ITS databases, we think that the bulk of species from floristic studies can, indeed, be identified with this method. We have to be aware that there are some taxonomic groups, which have too little ITS variability or too little amplification success, to be determined with this tool. A field, which could benefit from metabarcoding of eDNA, is community ecology of lichen-forming fungi, for example, understanding communities and species assemblages of lichenised fungi, their photobiont partners and other thallus-associated microorganisms. For example, species co-occurrences, based on eDNA, could be used to explore the concept of photobiont-mediated guilds (Rikkinen 2003). We have previously shown – based on the same DNA samples used here – that the communities of fungi, green algae and bacteria present on bark surfaces, strongly affect each other's beta diversities (Dreyling et al. 2024), suggesting that functional guilds, for example, of mycobionts and their photobiont partners, might be also be detected in the present data. Taxonomic assignments need to be carefully examined to assess if assignments are sensible for the geographic region of interest. Looking forward, the recent development of lichen specific databases might solve some of these issues. If eDNA biodiversity assessments are taken beyond the description of diversity, recently developed methods circumvent this issue altogether and are able to use unclassified taxa in the prediction of ecological states (Keck et al. 2023). Future studies of lichen biodiversity could employ these methods and expand the use of lichens as modern bio-monitoring agents.

Acknowledgements

We thank the managers of the three Exploratories, Max Müller, Julia Bass, Robert Künast, Anna K. Franke, Franca Marian and all former managers for their work in maintaining the plot and project infrastructure; Victoria Grießmeier for giving support through the central office, Andreas Ostrowski for managing the central data base and Markus Fischer, Eduard Linsenmair, Dominik Hesenmöller, Daniel Prati, Ingo Schöning, François Buscot, Ernst-Detlef Schulze, Wolfgang W. Weisser and the late Elisabeth Kalko for their role in setting up the Biodiversity Exploratories project. We thank the administration of the Hainich National Park, the UNESCO Biosphere Reserve Swabian Alb and the UNESCO Biosphere Reserve Schorfheide-Chorin, as well as all land owners for the excellent collaboration. Fieldwork permits were issued by the responsible state environmental offices of Baden-Württemberg, Thüringen and Brandenburg.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

The work has been (partly) funded by the DFG Priority Program 1374 “Infrastructure – Biodiversity- Exploratories” (SCHM 1711/8-1, FI 1246/6-1).

Author contributions

LD and IS conceptualized the method; LD collected eDNA samples and generated the taxonomic data; SB conducted the floristic mapping; LD, SB, HTL and IS curated the species list; LD compared the datasets; LD and IS wrote the manuscript; HTL and SB provided feedback.

Author ORCIDs

Lukas Dreyling  <https://orcid.org/0000-0001-9839-9504>

H. Thorsten Lumbsch  <https://orcid.org/0000-0003-1512-835X>

Imke Schmitt  <https://orcid.org/0000-0002-7381-0296>

Data availability

The raw sequencing data for this study is provided in the NCBI Sequence Read Archive under accession number SRR23371988. The code for the analysis is available at https://github.com/LukDrey/eDNA_lichen_survey.

References

Abarenkov K, Zirk A, Piirmann T, Pöhönen R, Ivanov F, Nilsson RH, Kõljalg U (2022) UNITE general FASTA release for eukaryotes. Version 27.10.2022. UNITE Community. <https://doi.org/10.15156/BIO/2483913>

- Abarenkov K, Nilsson RH, Larsson K-H, Taylor AFS, May TW, Frøslev TG, Pawlowska J, Lindahl B, Pöldmaa K, Truong C, Vu D, Hosoya T, Niskanen T, Piirmann T, Ivanov F, Zirk A, Peterson M, Cheeke TE, Ishigami Y, Jansson AT, Jeppesen TS, Kristiansson E, Mikryukov V, Miller JT, Oono R, Ossandon FJ, Paupério J, Saar I, Schigel D, Suija A, Tedersoo L, Kõljalg U (2023) The UNITE database for molecular identification and taxonomic communication of fungi and other eukaryotes: Sequences, taxa and classifications reconsidered. *Nucleic Acids Research*: gkad1039. <https://doi.org/10.1093/nar/gkad1039>
- Allen JL, Lendemer JC (2016) Climate change impacts on endemic, high-elevation lichens in a biodiversity hotspot. *Biodiversity and Conservation* 25(3): 555–568. <https://doi.org/10.1007/s10531-016-1071-4>
- Altermann S, Leavitt SD, Goward T, Nelsen MP, Lumbsch HT (2014) How Do You Solve a Problem like Letharia? A New Look at Cryptic Species in Lichen-Forming Fungi Using Bayesian Clustering and SNPs from Multilocus Sequence Data. *PLoS One* 9(5): e97556. <https://doi.org/10.1371/journal.pone.0097556>
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local alignment search tool. *Journal of Molecular Biology* 215(3): 403–410. [https://doi.org/10.1016/S0022-2836\(05\)80360-2](https://doi.org/10.1016/S0022-2836(05)80360-2)
- Aptroot A (2009) Chapter 23 – Lichens as an Indicator of Climate and Global Change. In: Letcher TM (Ed.) *Climate Change*. Elsevier, Amsterdam, 401–408. <https://doi.org/10.1016/B978-0-444-53301-2.00023-3>
- Asplund J, Wardle DA (2017) How lichens impact on terrestrial community and ecosystem properties. *Biological Reviews of the Cambridge Philosophical Society* 92(3): 1720–1738. <https://doi.org/10.1111/brv.12305>
- Asplund J, Larsson P, Vatne S, Gauslaa Y (2010) Gastropod grazing shapes the vertical distribution of epiphytic lichens in forest canopies. *Journal of Ecology* 98(1): 218–225. <https://doi.org/10.1111/j.1365-2745.2009.01584.x>
- Baldrian P (2017) Forest microbiome: Diversity, complexity and dynamics. *FEMS Microbiology Reviews* 41: 109–130. <https://doi.org/10.1093/femsre/fuw040>
- Bellemain E, Carlsen T, Brochmann C, Coissac E, Taberlet P, Kausserud H (2010) ITS as an environmental DNA barcode for fungi: An in silico approach reveals potential PCR biases. *BMC Microbiology* 10(1): 189. <https://doi.org/10.1186/1471-2180-10-189>
- Beng KC, Corlett RT (2020) Applications of environmental DNA (eDNA) in ecology and conservation: Opportunities, challenges and prospects. *Biodiversity and Conservation* 29(7): 2089–2121. <https://doi.org/10.1007/s10531-020-01980-0>
- Boch S, Prati D, Werth S, Rüetschi J, Fischer M (2011) Lichen Endozoochory by Snails. *PLoS One* 6(4): e18770. <https://doi.org/10.1371/journal.pone.0018770>
- Boch S, Prati D, Hessenmöller D, Schulze E-D, Fischer M (2013) Richness of Lichen Species, Especially of Threatened Ones, Is Promoted by Management Methods Furthering Stand Continuity. *PLoS One* 8(1): e55461. <https://doi.org/10.1371/journal.pone.0055461>
- Boch S, Saiz H, Allan E, Schall P, Prati D, Schulze E-D, Hessenmöller D, Sparrius LB, Fischer M (2021) Direct and Indirect Effects of Management Intensity and Environmental Factors on the Functional Diversity of Lichens in Central European Forests. *Microorganisms* 9(2): 463. <https://doi.org/10.3390/microorganisms9020463>
- Bradshaw M, Grewe F, Thomas A, Harrison CH, Lindgren H, Muggia L, St. Clair LL, Lumbsch HT, Leavitt SD (2020) Characterizing the ribosomal tandem repeat and its utility as a DNA barcode in lichen-forming fungi. *BMC Evolutionary Biology* 20: 2. <https://doi.org/10.1186/s12862-019-1571-4>

- Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP (2016) DADA2: High-resolution sample inference from Illumina amplicon data. *Nature Methods* 13(7): 581–583. <https://doi.org/10.1038/nmeth.3869>
- Camacho-Sanchez M, Burraco P, Gomez-Mestre I, Leonard JA (2013) Preservation of RNA and DNA from mammal samples under field conditions. *Molecular Ecology Resources* 13(4): 663–673. <https://doi.org/10.1111/1755-0998.12108>
- Campbell J, Fredeen AL, Prescott CE (2010) Decomposition and nutrient release from four epiphytic lichen litters in sub-boreal spruce forests. *Canadian Journal of Forest Research* 40(7): 1473–1484. <https://doi.org/10.1139/X10-071>
- Carter TS, Clark CM, Fenn ME, Jovan S, Perakis SS, Riddell J, Schaberg PG, Greaver TL, Hastings MG (2017) Mechanisms of nitrogen deposition effects on temperate forest lichens and trees. *Ecosphere* 8(3): e01717. <https://doi.org/10.1002/ecs2.1717>
- Coppins BJ, Coppins AM (2005) Lichens – The biodiversity value of western woodlands. *Botanical Journal of Scotland* 57(1–2): 141–153. <https://doi.org/10.1080/03746600508685093>
- Cordier T, Alonso-Sáez L, Apothéloz-Perret-Gentil L, Aylagas E, Bohan DA, Bouchez A, Chariton A, Creer S, Frühe L, Keck F, Keeley N, Laroche O, Leese F, Pochon X, Stoeck T, Pawlowski J, Lanzén A (2021) Ecosystems monitoring powered by environmental genomics: A review of current strategies with an implementation roadmap. *Molecular Ecology* 30(13): 2937–2958. <https://doi.org/10.1111/mec.15472>
- Crespo A, Lumbsch HT (2010) Cryptic species in lichen-forming fungi. *IMA Fungus* 1(2): 167–170. <https://doi.org/10.5598/imafungus.2010.01.02.09>
- Crous PW, Gams W, Stalpers JA, Robert V, Stegehuis G (2004) MycoBank: An online initiative to launch mycology into the 21st century. *Studies in Mycology* 50: 19–22.
- Czarnota P, Guzow-Krzemińska B (2018) *Bacidina mendax* sp. nov., a new widespread species in Central Europe, together with a new combination within the genus *Bacidina*. *Lichenologist* (London, England) 50(1): 43–57. <https://doi.org/10.1017/S0024282917000627>
- Deagle BE, Eveson JP, Jarman SN (2006) Quantification of damage in DNA recovered from highly degraded samples – a case study on DNA in faeces. *Frontiers in Zoology* 3(1): 11. <https://doi.org/10.1186/1742-9994-3-11>
- Deiner K, Bik HM, Mächler E, Seymour M, Lacoursière-Roussel A, Altermatt F, Creer S, Bista I, Lodge DM, de Vere N, Pfrender ME, Bernatchez L (2017) Environmental DNA metabarcoding: Transforming how we survey animal and plant communities. *Molecular Ecology* 26(21): 5872–5895. <https://doi.org/10.1111/mec.14350>
- Divakar PK, Leavitt SD, Molina MC, Del-Prado R, Lumbsch HT, Crespo A (2016) A DNA barcoding approach for identification of hidden diversity in Parmeliaceae (Ascomycota): *Parmelia sensu stricto* as a case study. *Botanical Journal of the Linnean Society* 180(1): 21–29. <https://doi.org/10.1111/boj.12358>
- Dreyling L, Schmitt I, Dal Grande F (2022) Tree Size Drives Diversity and Community Structure of Microbial Communities on the Bark of Beech (*Fagus sylvatica*). *Frontiers in Forests and Global Change* 5. <https://doi.org/10.3389/ffgc.2022.858382> [October 4, 2023]
- Dreyling L, Penone C, Schenk NV, Schmitt I, Dal Grande F (2024) Biotic interactions outweigh abiotic factors as drivers of bark microbial communities in central European forests. *ISME Communications* 4(1): ycae012. <https://doi.org/10.1093/ismeco/ycae012>
- Dymytrova L (2011) Notes on the genus *Scoliciosporum* (Lecanorales, Ascomycota) in Ukraine. *Polish Botanical Journal* 56: 61–75.

- Ellis CJ (2012) Lichen epiphyte diversity: A species, community and trait-based review. *Perspectives in Plant Ecology, Evolution and Systematics* 14(2): 131–152. <https://doi.org/10.1016/j.ppees.2011.10.001>
- Farkas E, Varga N, Veres K, Matus G, Sinigla M, Lőkös L (2022) Distribution Types of Lichens in Hungary That Indicate Changing Environmental Conditions. *Journal of Fungi (Basel, Switzerland)* 8(6): 600. <https://doi.org/10.3390/jof8060600>
- Fediajevaite J, Priestley V, Arnold R, Savolainen V (2021) Meta-analysis shows that environmental DNA outperforms traditional surveys, but warrants better reporting standards. *Ecology and Evolution* 11(9): 4803–4815. <https://doi.org/10.1002/ece3.7382>
- Fischer M, Bossdorf O, Gockel S, Hänsel F, Hemp A, Hessenmöller D, Korte G, Nieschulze J, Pfeiffer S, Prati D, Renner S, Schöning I, Schumacher U, Wells K, Buscot F, Kalko EKV, Linsenmair KE, Schulze E-D, Weisser WW (2010) Implementing large-scale and long-term functional biodiversity research: The Biodiversity Exploratories. *Basic and Applied Ecology* 11(6): 473–485. <https://doi.org/10.1016/j.baae.2010.07.009>
- Fрати L, Brunialti G (2023) Recent Trends and Future Challenges for Lichen Biomonitoring in Forests. *Forests* 14(3): 647. <https://doi.org/10.3390/f14030647>
- Frøslev TG, Kjølner R, Bruun HH, Ejrnæs R, Hansen AJ, Læssøe T, Heilmann-Clausen J (2019) Man against machine: Do fungal fruitbodies and eDNA give similar biodiversity assessments across broad environmental gradients? *Biological Conservation* 233: 201–212. <https://doi.org/10.1016/j.biocon.2019.02.038>
- Gadsdon SR, Dagley JR, Wolseley PA, Power SA (2010) Relationships between lichen community composition and concentrations of NO₂ and NH₃. *Environmental Pollution* 158(8): 2553–2560. <https://doi.org/10.1016/j.envpol.2010.05.019>
- Gilbert OL (1992) Lichen reinvasion with declining air pollution. In: Bates JW, Farmer AM (Eds) *Bryophytes and Lichens in a Changing Environment*. Oxford University Press. <https://doi.org/10.1093/oso/9780198542919.003.0006>
- Giordani P, Brunialti G, Benesperi R, Rizzi G, Frati L, Modenesi P (2009) Rapid biodiversity assessment in lichen diversity surveys: Implications for quality assurance. *Journal of Environmental Monitoring* 11(4): 730–735. <https://doi.org/10.1039/b818173j>
- Gloor GB, Macklaim JM, Pawlowsky-Glahn V, Egozcue JJ (2017) Microbiome Datasets Are Compositional: And This Is Not Optional. *Frontiers in Microbiology* 8: 2224. <https://doi.org/10.3389/fmicb.2017.02224>
- Gueidan C, Li L (2022) A long-read amplicon approach to scaling up the metabarcoding of lichen herbarium specimens. *MycKeys* 86: 195–212. <https://doi.org/10.3897/mycokeys.86.77431>
- Henrie JR, Thomson BM, Yungfleisch AA, Kerr M, Leavitt SD (2022) Characterizing Crustose Lichen Communities – DNA Metabarcoding Reveals More than Meets the Eye. *Diversity* 14(9): 766. <https://doi.org/10.3390/d14090766>
- Hofmann B, Dreyling L, Dal Grande F, Otte J, Schmitt I (2023) Habitat and tree species identity shape aboveground and belowground fungal communities in central European forests. *Frontiers in Microbiology* 14: 1067906. <https://doi.org/10.3389/fmicb.2023.1067906>
- Hofmeister J, Hošek J, Malíček J, Palice Z, Syrovátková L, Steinová J, Černajová I (2016) Large beech (*Fagus sylvatica*) trees as 'lifeboats' for lichen diversity in central European forests. *Biodiversity and Conservation* 25(6): 1073–1090. <https://doi.org/10.1007/s10531-016-1106-x>
- Hultengren S, Gralén H, Pleijel H (2004) Recovery of the Epiphytic Lichen Flora Following Air Quality Improvement in South-West Sweden. *Water, Air, and Soil Pollution* 154(1–4): 203–211. <https://doi.org/10.1023/B:WATE.0000022967.35036.ca>

- Ihrmark K, Bödeker ITM, Cruz-Martinez K, Friberg H, Kubartova A, Schenck J, Strid Y, Stenlid J, Brandström-Durling M, Clemmensen KE, Lindahl BD (2012) New primers to amplify the fungal ITS2 region – evaluation by 454-sequencing of artificial and natural communities. *FEMS Microbiology Ecology* 82(3): 666–677. <https://doi.org/10.1111/j.1574-6941.2012.01437.x>
- Jüriado I, Paal J, Liira J (2003) Epiphytic and epixylic lichen species diversity in Estonian natural forests. *Biodiversity and Conservation* 12(8): 1587–1607. <https://doi.org/10.1023/A:1023645730446>
- Keck F, Blackman RC, Bossart R, Brantschen J, Couton M, Hürlemann S, Kirschner D, Locher N, Zhang H, Altermatt F (2022) Meta-analysis shows both congruence and complementarity of DNA and eDNA metabarcoding to traditional methods for biological community assessment. *Molecular Ecology* 31(6): 1820–1835. <https://doi.org/10.1111/mec.16364>
- Keck F, Brantschen J, Altermatt F (2023) A combination of machine-learning and eDNA reveals the genetic signature of environmental change at the landscape levels. *Molecular Ecology* 32(17): 4791–4800. <https://doi.org/10.1111/mec.17073>
- Kelly LJ, Hollingsworth PM, Coppins BJ, Ellis CJ, Harrold P, Tosh J, Yahr R (2011) DNA barcoding of lichenized fungi demonstrates high identification success in a floristic context. *The New Phytologist* 191(1): 288–300. <https://doi.org/10.1111/j.1469-8137.2011.03677.x>
- Kerr M, Leavitt SD (2023) A Custom Regional DNA Barcode Reference Library for Lichen-Forming Fungi of the Intermountain West, USA, Increases Successful Specimen Identification. *Journal of Fungi (Basel, Switzerland)* 9(7): 741. <https://doi.org/10.3390/jof9070741>
- Klaus VH, Hölzel N, Boch S, Müller J, Socher AS, Prati D, Fischer M, Kleinebecker T (2013) Direct and indirect associations between plant species richness and productivity in grasslands: Regional differences preclude simple generalization of productivity-biodiversity relationships. *Preslia* 85: 97–112. <https://doi.org/10.7892/boris.38709>
- Knops JMH, Iii THN, Boucher VL, Schlesinger WH (1991) Mineral Cycling and Epiphytic Lichens: Implications at the Ecosystem Level. *Lichenologist (London, England)* 23(3): 309–321. <https://doi.org/10.1017/S0024282991000452>
- Knops JMH, Nash TH III, Schlesinger WH (1996) The Influence of Epiphytic Lichens on the Nutrient Cycling of an Oak Woodland. *Ecological Monographs* 66(2): 159–179. <https://doi.org/10.2307/2963473>
- Kowalewska A, Kukwa M (2003) Additions to the Polish lichen flora. *Graphis Scripta*.
- Kraichak E, Allende L, Obermayer W, Lücking R, Lumbsch HT (2019) Scale-dependent co-occurrence patterns of closely related genotypes in a lichen species complex. *Plant and Fungal Systematics* 64(2): 163–172. <https://doi.org/10.2478/pfs-2019-0017>
- Kranner I, Zorn M, Turk B, Wornik S, Beckett RP, Batič F (2003) Biochemical traits of lichens differing in relative desiccation tolerance. *The New Phytologist* 160(1): 167–176. <https://doi.org/10.1046/j.1469-8137.2003.00852.x>
- Lackovičová A, Guttová A (2005) Genus *Dimerella* (Coenogoniaceae, lichenized Ascomycota) in Slovakia. *Acta Botanica Croatica* 64: 289–301.
- Larsen HME, Hansen ES, Nord-Larsen T, Rasmussen HN (2020) Deciduous trees as lichen phorophytes: Biodiversity and colonization patterns under common garden conditions. *Lichenologist (London, England)* 52(3): 221–232. <https://doi.org/10.1017/S0024282920000018>

- Launis A, Pykälä J, Boom PVD, Sérusiaux E, Myllys L (2019) Four new epiphytic species in the *Micarea prasina* group from Europe. *Lichenologist* (London, England) 51(1): 7–25. <https://doi.org/10.1017/S0024282918000555>
- Liu D, Wang L, Wang XY, Hur J-S (2019) Two New Species of the Genus *Candelariella* from China and Korea. *Mycobiology* 47(1): 40–49. <https://doi.org/10.1080/12298093.2019.1583785>
- Löhmus P, Löhmus A (2019) The Potential of Production Forests for Sustaining Lichen Diversity: A Perspective on Sustainable Forest Management. *Forests* 10(12): 1063. <https://doi.org/10.3390/f10121063>
- Lubek A, Kukwa M, Jaroszewicz B, Czortek P (2021) Shifts in Lichen Species and Functional Diversity in a Primeval Forest Ecosystem as a Response to Environmental Changes. *Forests* 12(6): 686. <https://doi.org/10.3390/f12060686>
- Lücking R, Aime MC, Robbertse B, Miller AN, Ariyawansa HA, Aoki T, Cardinali G, Crous PW, Druzhinina IS, Geiser DM, Hawksworth DL, Hyde KD, Irinyi L, Jeewon R, Johnston PR, Kirk PM, Malosso E, May TW, Meyer W, Öpik M, Robert V, Stadler M, Thines M, Vu D, Yurkov AM, Zhang N, Schoch CL (2020) Unambiguous identification of fungi: Where do we stand and how accurate and precise is fungal DNA barcoding? *IMA Fungus* 11(1): 14. <https://doi.org/10.1186/s43008-020-00033-z>
- Marini L, Nascimbene J, Nimis PL (2011) Large-scale patterns of epiphytic lichen species richness: Photobiont-dependent response to climate and forest structure. *The Science of the Total Environment* 409(20): 4381–4386. <https://doi.org/10.1016/j.scitotenv.2011.07.010>
- Marthinsen G, Rui S, Timdal E (2019) OLICH: A reference library of DNA barcodes for Nordic lichens. *Biodiversity Data Journal* 7: e36252. <https://doi.org/10.3897/BDJ.7.e36252>
- Martin M (2011) Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.Journal* 17(1): 10. <https://doi.org/10.14806/ej.17.1.200>
- Miller JED, Root HT, Safford HD (2018) Altered fire regimes cause long-term lichen diversity losses. *Global Change Biology* 24(10): 4909–4918. <https://doi.org/10.1111/gcb.14393>
- Nascimbene J, Thor G, Nimis PL (2013) Effects of forest management on epiphytic lichens in temperate deciduous forests of Europe – A review. *Forest Ecology and Management* 298: 27–38. <https://doi.org/10.1016/j.foreco.2013.03.008>
- Nascimbene J, Casazza G, Benesperi R, Catalano I, Cataldo D, Grillo M, Isocrono D, Matteucci E, Ongaro S, Potenza G, Puntillo D, Ravera S, Zedda L, Giordani P (2016) Climate change fosters the decline of epiphytic *Lobaria* species in Italy. *Biological Conservation* 201: 377–384. <https://doi.org/10.1016/j.biocon.2016.08.003>
- Nash TH III, Gries C (2002) Lichens as Bioindicators of Sulfur Dioxide. *Symbiosis*: 1–21.
- Nelsen MP, Lumbsch HT (2020) A data-driven evaluation of lichen climate change indicators in Central Europe. *Biodiversity and Conservation* 29(14): 3959–3971. <https://doi.org/10.1007/s10531-020-02057-8>
- Nguyen NH, Song Z, Bates ST, Branco S, Tedersoo L, Menke J, Schilling JS, Kennedy PG (2016) FUNGuild: An open annotation tool for parsing fungal community datasets by ecological guild. *Fungal Ecology* 20: 241–248. <https://doi.org/10.1016/j.funeco.2015.06.006>
- Pike LH (1978) The Importance of Epiphytic Lichens in Mineral Cycling. *The Bryologist* 81(2): 247–257. <https://doi.org/10.2307/3242186>
- Pinho P, Augusto S, Martins-Loução MA, Pereira MJ, Soares A, Máguas C, Branquinho C (2008) Causes of change in nitrophytic and oligotrophic lichen species in a Medi-

- terranean climate: Impact of land cover and atmospheric pollutants. *Environmental Pollution* 154(3): 380–389. <https://doi.org/10.1016/j.envpol.2007.11.028>
- Pino-Bodas R, Martín MP, Burgaz AR, Lumbsch HT (2013) Species delimitation in *Cladonia* (Ascomycota): A challenge to the DNA barcoding philosophy. *Molecular Ecology Resources* 13(6): 1058–1068. <https://doi.org/10.1111/1755-0998.12086>
- Purvis OW, Chimonides J, Din V, Erotokritou L, Jeffries T, Jones GC, Louwhoff S, Read H, Spiro B (2003) Which factors are responsible for the changing lichen floras of London? *The Science of the Total Environment* 310(1–3): 179–189. [https://doi.org/10.1016/S0048-9697\(02\)00638-1](https://doi.org/10.1016/S0048-9697(02)00638-1)
- Reiners WA, Olson RK (1984) Effects of canopy components on throughfall chemistry: An experimental analysis. *Oecologia* 63(3): 320–330. <https://doi.org/10.1007/BF00390660>
- Rikkinen J (2003) Ecological and evolutionary role of photobiont-mediated guilds in lichens. *Symbiosis* 2003: 443–450.
- Rivas Plata E, Parnmen S, Staiger B, Mangold A, Frisch A, Weerakoon G, Hernandez J, Caceres M, Kalb K, Sipman H, Common R, Nelsen M, Luecking R, Lumbsch T (2013) A molecular phylogeny of Graphidaceae (Ascomycota, Lecanoromycetes, Ostropales) including 428 species. *MycKeys* 6: 55–94. <https://doi.org/10.3897/mycokeys.6.3482>
- Robert V, Vu D, Amor ABH, van de Wiele N, Brouwer C, Jabas B, Szoke S, Dridi A, Triki M, Daoud S, Chouchen O, Vaas L, de Cock A, Stalpers JA, Stalpers D, Verkley GJM, Groenewald M, dos Santos FB, Stegehuis G, Li W, Wu L, Zhang R, Ma J, Zhou M, Gorjón SP, Eurwilaichitr L, Ingsriswang S, Hansen K, Schoch C, Robbertse B, Irinyi L, Meyer W, Cardinali G, Hawksworth DL, Taylor JW, Crous PW (2013) MycoBank gearing up for new horizons. *IMA Fungus* 4(2): 371–379. <https://doi.org/10.5598/ima fungus.2013.04.02.16>
- Rose CI, Hawksworth DL (1981) Lichen recolonization in London's cleaner air. *Nature* 289(5795): 289–292. <https://doi.org/10.1038/289289a0>
- Sayers EW, Bolton EE, Brister JR, Canese K, Chan J, Comeau DC, Connor R, Funk K, Kelly C, Kim S, Madej T, Marchler-Bauer A, Lanczycki C, Lathrop S, Lu Z, Thibaud-Nissen F, Murphy T, Phan L, Skripchenko Y, Tse T, Wang J, Williams R, Trawick BW, Pruitt KD, Sherry ST (2022) Database resources of the national center for biotechnology information. *Nucleic Acids Research* 50(D1): D20–D26. <https://doi.org/10.1093/nar/gkab1112>
- Schall P, Ammer C (2018) Stand composition based on 2nd forest inventory (abundance, basal area, crown projection area and volume) on all forest EPs, 2014 – 2018. Version 2. Biodiversity Exploratories Information System. Dataset. <https://www.bexis.uni-jena.de/ddm/data/Showdata/22907>
- Schmitt I, Lumbsch HT (2004) Molecular phylogeny of the Pertusariaceae supports secondary chemistry as an important systematic character set in lichen-forming ascomycetes. *Molecular Phylogenetics and Evolution* 33(1): 43–55. <https://doi.org/10.1016/j.ympev.2004.04.014>
- Schoch CL, Seifert KA, Huhndorf S, Robert V, Spouge JL, Levesque CA, Chen W, et al. (2012) Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for Fungi. *Proceedings of the National Academy of Sciences* 109: 6241–6246. <https://doi.org/10.1073/pnas.1117018109>
- Shirouzu T, Uno K, Hosaka K, Hosoya T (2016) Early-diverging wood-decaying fungi detected using three complementary sampling methods. *Molecular Phylogenetics and Evolution* 98: 11–20. <https://doi.org/10.1016/j.ympev.2016.01.015>

- Simon DM, Hummel CL, Sheeley SL, Bhattacharya D (2005) Heterogeneity of intron presence or absence in rDNA genes of the lichen species *Physcia aipolia* and *P. stellaris*. *Current Genetics* 47(6): 389–399. <https://doi.org/10.1007/s00294-005-0581-5>
- Simons NK, Gossner MM, Lewinsohn TM, Boch S, Lange M, Müller J, Pašalić E, Socher SA, Türke M, Fischer M, Weisser WW (2014) Resource-Mediated Indirect Effects of Grassland Management on Arthropod Diversity. *PLoS ONE* 9: e107033. <https://doi.org/10.1371/journal.pone.0107033>
- Taberlet P, Coissac E, Pompanon F, Brochmann C, Willerslev E (2012) Towards next-generation biodiversity assessment using DNA metabarcoding. *Molecular Ecology* 21(8): 2045–2050. <https://doi.org/10.1111/j.1365-294X.2012.05470.x>
- Umweltbundesamt (2005) 8 Daten zur Umwelt. 8: 2005. Berlin, 352 pp.
- van Herk CM, Aptroot A, van Dobben HF (2002) Long-Term Monitoring in the Netherlands Suggests that Lichens Respond to Global Warming. *Lichenologist* (London, England) 34(2): 141–154. <https://doi.org/10.1006/lich.2002.0378>
- Van Stan JT II, Pypker TG (2015) A review and evaluation of forest canopy epiphyte roles in the partitioning and chemical alteration of precipitation. *The Science of the Total Environment* 536: 813–824. <https://doi.org/10.1016/j.scitotenv.2015.07.134>
- Větrovský T, Morais D, Kohout P, Lepinay C, Algora C, Awokunle Hollá S, Bahnmann BD, Bílohnědá K, Brabcová V, D'Alò F, Human ZR, Jomura M, Kolařík M, Kvasničková J, Lladó S, López-Mondéjar R, Martinović T, Mašínová T, Meszárosová L, Michalčíková L, Michalová T, Mundra S, Navrátilová D, Odriozola I, Piché-Choquette S, Štursová M, Švec K, Tláškal V, Urbanová M, Vlk L, Voříšková J, Žifčáková L, Baldrian P (2020) GlobalFungi, a global database of fungal occurrences from high-throughput-sequencing metabarcoding studies. *Scientific Data* 7(1): 228. <https://doi.org/10.1038/s41597-020-0567-7>
- Vondrák J, Malíček J, Palice Z, Coppins B, Kukwa M, Czarnota P, Sanderson N, Acton A (2016) Methods for obtaining more complete species lists in surveys of lichen biodiversity. *Nordic Journal of Botany* 34(5): 619–626. <https://doi.org/10.1111/njb.01053>
- Vondrák J, Svoboda S, Košnar J, Malíček J, Šoun J, Frolov I, Svensson M, Novotný P, Palice Z (2023) Martin7: A reference database of DNA barcodes for European epiphytic lichens and its taxonomic implications. *Preslia* 95(3): 311–345. <https://doi.org/10.23855/preslia.2023.311>
- White TJ, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal rna genes for phylogenetics. In: Innis M, Gelfand DH, Sninsky JJ, White TJ (Eds) *PCR Protocols*. Academic Press, San Diego, CA, 315–322. <https://doi.org/10.1016/B978-0-12-372180-8.50042-1>
- Wirth V, Hauck M, Schultz M (2013) *Die Flechten Deutschlands*. Ulmer, Stuttgart, 2 pp.
- Wright B, St. Clair LL, Leavitt SD (2019) Is targeted community DNA metabarcoding suitable for biodiversity inventories of lichen-forming fungi? *Ecological Indicators* 98: 812–820. <https://doi.org/10.1016/j.ecolind.2018.11.061>
- Ye J, Coulouris G, Zaretskaya I, Cutcutache I, Rozen S, Madden TL (2012) Primer-BLAST: A tool to design target-specific primers for polymerase chain reaction. *BMC Bioinformatics* 13(1): 134. <https://doi.org/10.1186/1471-2105-13-134>
- Yoccoz NG (2012) The future of environmental DNA in ecology. *Molecular Ecology* 21(8): 2031–2038. <https://doi.org/10.1111/j.1365-294X.2012.05505.x>

Supplementary material 1

List of all species found in this study

Authors: Lukas Dreyling, Steffen Boch, H. Thorsten Lumbsch, Imke Schmitt

Data type: csv

Explanation note: The table contains information on whether the species was identified with either the eDNA method or the floristic survey and in how many plots it occurred.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/mycokeys.106.117540.suppl1>